

Fourier, Gegenbauer and Jacobi expansions for a power-law fundamental solution of the polyharmonic equation and polyspherical addition theorems

H S Cohl¹

¹Information Technology Laboratory, National Institute of Standards and Technology, Gaithersburg, MD, USA

E-mail: hcohl@nist.gov

Abstract. We develop complex Jacobi, Gegenbauer and Chebyshev polynomial expansions for the kernels associated with power-law fundamental solutions of the polyharmonic equation on d -dimensional Euclidean space. From these series representations we derive Fourier expansions in rotationally-invariant coordinate systems and Gegenbauer polynomial expansions in Vilenkin's polyspherical coordinates. The resulting Fourier and Gegenbauer coefficients are given in terms of associated Legendre functions. We compare both of these expansions to generate addition theorems for the azimuthal Fourier coefficients.

PACS numbers: 02.30.Em, 02.30.Gp, 02.30.Jr, 02.30.Nw, 02.40.Ky, 02.40.Vh

AMS classification scheme numbers: 35A08, 31B30, 31C12, 33C05, 42A16

1. Introduction

We have developed a technique for constructing addition theorems for the azimuthal Fourier coefficients of fundamental solutions for linear homogenous partial differential equations on d -dimensional Riemannian manifolds. In our technique, one must construct azimuthal Fourier expansions for this fundamental solution and compare it with eigenfunction expansions for that fundamental solution in particular separable rotationally-invariant coordinate systems. The construction of eigenfunction expansions for fundamental solutions in separable coordinate systems is in general non-trivial.

In this paper, we apply this technique to generate addition theorems from eigenfunction expansions for a fundamental solution of the polyharmonic operator on d -dimensional Euclidean space $\mathbf{R}^d$ in Vilenkin-Kuznetsov-Smorodinskii polyspherical coordinate systems [25, 26] (hereafter Vilenkin). These coordinate systems are subgroup type coordinate systems, as they can be described using a maximal subgroup chain, and are perhaps the most studied rotationally-invariant coordinate systems in existence for this operator. Here we have computed azimuthal Fourier expansions for a fundamental solution of this operator, as well as the corresponding eigenfunction expansions in polyspherical coordinate systems. In each case, the comparison of these two expansions yields new addition theorems for the azimuthal Fourier coefficients.

A fundamental solution of the polyharmonic equation on $\mathbf{R}^d$ is given either in terms of solutions which have power-law or logarithmic behavior as a function of the distance between two points on this space (see for instance Schwartz (1950) ([22, p. 45]); Gel'fand & Shilov (1964) ([12], p. 202)). In a recent paper (Cohl & Dominici (2010) [5]), we derived a complex identity which determined the Fourier coefficients of a power-law fundamental solution of the polyharmonic equation. The Fourier coefficients were seen to be given in terms of associated Legendre functions. The eigenfunction expansions for a power-law fundamental solution of the polyharmonic equation in a particular polyspherical coordinate system can be derived using a Gegenbauer polynomial expansion for the relevant kernel (see Corollary 5.4 below) in conjunction with the addition theorem for hyperspherical harmonics. This addition theorem is given by

$$C_n^{d/2-1}(\cos \gamma) = \frac{2(d-2)\pi^{d/2}}{(2n+d-2)\Gamma(d/2)} \sum_K Y_n^K(\hat{\mathbf{x}}) \overline{Y_n^K(\hat{\mathbf{x}}')}, \quad (1)$$

(for a proof see Wen & Avery (1985) [27]; see also section 10.2.1 in Fano & Rau (1996) [11]) where K stands for a set of $(d-2)$ -quantum numbers identifying degenerate harmonics for a given value of $n \in \mathbf{N}_0$, and $\cos \gamma$ is the cosine of the separation angle between two arbitrary vectors $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$ (see (12) below). The functions $Y_n^K : \mathbf{S}^{d-1} \rightarrow \mathbf{C}$ are the normalized hyperspherical harmonics, with $\hat{\mathbf{x}}, \hat{\mathbf{x}}' \in \mathbf{S}^{d-1}$ being unit vectors in the direction of $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$ respectively. The Riemannian manifold $\mathbf{S}_R^d$ is defined as the set of all points in $\mathbf{R}^{d+1}$ such that $x_0^2 + \dots + x_d^2 = R^2$ ($R > 0$), with the metric induced from that of the ambient Euclidean space. We denote the d -dimensional hypersphere of unit radius by $\mathbf{S}^d := \mathbf{S}_1^d$.

There will be a separate addition theorem corresponding to each rotationally-invariant coordinate system. It should be mentioned that the large number and complexity of coordinate systems increases multi-fold when one increases dimension. One could also allow for the possibility of non-subgroup type rotationally-invariant coordinate systems such as parabolic and spheroidal coordinates but these will not be addressed in the present paper. See for instance Kalnins (1986) [17] and Cohl *et al.* (2000) [8] for some examples of this type of coordinate systems connected with the Laplace operator on $\mathbf{R}^3$. In this paper we give examples of addition theorems from a power-law fundamental solution of the polyharmonic operator in Euclidean space using Vilenkin's polyspherical coordinate systems in low dimensions (i.e., $d \in \{3, 4\}$) and also in higher dimensions.

This paper is organized as follows. In section 2 we describe the kernels associated with a fundamental solution of the polyharmonic equation on Euclidean space $\mathbf{R}^d$. In section 3 we describe Vilenkin's polyspherical coordinate systems. In section 4 we describe the normalized hyperspherical harmonics associated with Vilenkin's polyspherical coordinates. In section 5 we prove several theorems associated with Jacobi, Gegenbauer, and Chebyshev polynomial expansions of the kernels associated with power-law fundamental solutions of the polyharmonic equation on Euclidean space. In section 6 we derive and discuss addition theorems in Vilenkin's polyspherical coordinates for the Fourier coefficients of a fundamental solution for the polyharmonic equation on $\mathbf{R}^d$.

Throughout this paper we rely on the following definitions. Let $a_1, a_2, a_3, \dots \in \mathbf{C}$, with $\mathbf{C}$ being the set of complex numbers. If $i, j \in \mathbf{Z}$ and $j < i$, then $\sum_{n=i}^j a_n = 0$ and $\prod_{n=i}^j a_n = 1$. The set of natural numbers is given by $\mathbf{N} := \{1, 2, 3, \dots\}$, the set $\mathbf{N}_0 := \{0, 1, 2, \dots\} = \mathbf{N} \cup \{0\}$, and $\mathbf{Z} := \{0, \pm 1, \pm 2, \dots\}$. The sets $\mathbf{Q}$ and $\mathbf{R}$ represents the rational and real numbers respectively.

2. Fundamental solution of the polyharmonic equation and the non-logarithmic Fourier series

If $\Phi : \mathbf{R}^d \rightarrow \mathbf{R}$ satisfies the polyharmonic equation given by

$$(-\Delta)^k \Phi(\mathbf{x}) = 0, \quad (2)$$

where $\mathbf{x} \in \mathbf{R}^d$, $\Delta : C^p(\mathbf{R}^d) \rightarrow C^{p-2}(\mathbf{R}^d)$ for $p \geq 2$, is the Laplacian operator defined by

$$\Delta := \frac{\partial^2}{\partial x_1^2} + \dots + \frac{\partial^2}{\partial x_d^2},$$

where $k \in \mathbf{N}$ and $\Phi \in C^{2k}(\mathbf{R}^d)$, then Φ is called polyharmonic. The inhomogeneous polyharmonic equation is given by

$$(-\Delta)^k \Phi(\mathbf{x}) = \rho(\mathbf{x}), \quad (3)$$

where we take ρ to be an integrable function so that a solution to (3) exists. A fundamental solution for the polyharmonic equation on $\mathbf{R}^d$ is a function $\mathcal{G}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \rightarrow \mathbf{R}$ which satisfies the equation

$$(-\Delta)^k \mathcal{G}_k^d(\mathbf{x}, \mathbf{x}') = \delta(\mathbf{x} - \mathbf{x}'), \quad (4)$$

where δ is the Dirac delta function (generalized function) and $\mathbf{x}' \in \mathbf{R}^d$. The Euclidean inner product $(\cdot, \cdot) : \mathbf{R}^d \times \mathbf{R}^d \rightarrow \mathbf{R}$ defined by

$$(\mathbf{x}, \mathbf{x}') := x_1 x'_1 + \dots + x_d x'_d, \quad (5)$$

induces a norm (the Euclidean norm) $\|\cdot\| : \mathbf{R}^d \rightarrow [0, \infty)$, on the finite-dimensional vector space $\mathbf{R}^d$, given by $\|\mathbf{x}\| := \sqrt{(\mathbf{x}, \mathbf{x})}$.

In the rest of this paper, we will use the gamma function $\Gamma : \mathbf{C} \setminus -\mathbf{N}_0 \rightarrow \mathbf{C}$, which is a natural generalization of the factorial function (see for instance Chapter 5 in Olver *et al.* (2010) [21]). A fundamental solution of the polyharmonic equation is given by the following theorem.

Theorem 2.1. *Let $d, k \in \mathbf{N}$. Define $\mathcal{G}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \rightarrow \mathbf{R}$ such that*

$$\mathcal{G}_k^d(\mathbf{x}, \mathbf{x}') := \begin{cases} \frac{(-1)^{k+d/2+1} \|\mathbf{x} - \mathbf{x}'\|^{2k-d}}{(k-1)! (k-d/2)! 2^{2k-1} \pi^{d/2}} (\log \|\mathbf{x} - \mathbf{x}'\| - \beta_{k-d/2,d}) & \text{if } d \text{ even, } k \geq d/2, \\ \frac{\Gamma(d/2 - k) \|\mathbf{x} - \mathbf{x}'\|^{2k-d}}{(k-1)! 2^{2k} \pi^{d/2}} & \text{otherwise,} \end{cases}$$

where $\beta_{p,d} \in \mathbf{Q}$ is defined as $\beta_{p,d} := \frac{1}{2} [H_p + H_{d/2+p-1} - H_{d/2-1}]$, with H_j being the j th harmonic number

$$H_j := \sum_{i=1}^j \frac{1}{i},$$

then $\mathcal{G}_k^d$ is a normalized fundamental solution for $(-\Delta)^k$ on $\mathbf{R}^d$.

Proof. See Cohl (2010) [4] and Boyling (1996) [3].

We consider parametrizations of points on $\mathbf{R}^d$ which are given in terms of coordinate systems whose coordinates are curvilinear, i.e., based on some transformation which converts the Cartesian coordinates to a coordinate system with the same number of coordinates in which the coordinate lines are curved. We consider solutions of the polyharmonic equation (2), in a curvilinear coordinate system, which arises through the theory of separation of variables. We refer to coordinate systems which yield solutions through the separation of variables method as separable. In this paper we restrict our attention to separable rotationally-invariant coordinate systems for the polyharmonic equation on $\mathbf{R}^d$ which are given by

$$\left. \begin{aligned} x_1 &= R(\xi_1, \dots, \xi_{d-1}) \cos \phi \\ x_2 &= R(\xi_1, \dots, \xi_{d-1}) \sin \phi \\ x_3 &= x_3(\xi_1, \dots, \xi_{d-1}) \\ &\vdots \\ x_d &= x_d(\xi_1, \dots, \xi_{d-1}) \end{aligned} \right\}. \quad (6)$$

These coordinate systems are described by d coordinates: an angle $\phi \in [0, 2\pi)$ plus $(d-1)$ -curvilinear coordinates $(\xi_1, \dots, \xi_{d-1})$. Rotationally-invariant coordinate systems

parametrize points on the $(d - 1)$ -dimensional half-hyperplane given by $\phi = \text{const}$ and $R \geq 0$ using the curvilinear coordinates $(\xi_1, \dots, \xi_{d-1})$. A separable rotationally-invariant coordinate system transforms the polyharmonic equation into a set of d -uncoupled ordinary differential equations with separation constants $m \in \mathbf{Z}$ and $k_j \in \mathbf{R}$ for $1 \leq j \leq d - 2$. For a separable rotationally-invariant coordinate system, this uncoupling is accomplished, in general, by assuming a solution to (2) of the form

$$\Phi(x) = e^{im\phi} \mathcal{R}(\xi_1, \dots, \xi_{d-1}) \prod_{i=1}^{d-1} A_i(\xi_i, m, k_1, \dots, k_{d-2}),$$

where the properties of the functions $\mathcal{R}$ and A_i , for $1 \leq i \leq d - 1$, and the constants k_j for $1 \leq j \leq d - 2$, depend on the specific separable rotationally-invariant coordinate system in question. Separable coordinate systems are divided into two separate classes, those which are simply separable ($\mathcal{R} = \text{const}$), and those which are $\mathcal{R}$ -separable. For a general description of the theory of separation of variables see Miller (1977) [20].

The Euclidean distance between two points $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$, expressed in a rotationally-invariant coordinate system, is given by

$$\|\mathbf{x} - \mathbf{x}'\| = \sqrt{2RR'} [\chi - \cos(\phi - \phi')]^{1/2}, \quad (7)$$

where the toroidal parameter $\chi > 1$, is given by

$$\chi := \frac{R^2 + R'^2 + \sum_{i=3}^d (x_i - x'_i)^2}{2RR'}, \quad (8)$$

where $R, R' \in (0, \infty)$ are defined in (6) for $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$. The hypersurfaces given by $\chi > 1$ equals constant are independent of coordinate system and represent hypertori of revolution.

From Theorem 2.1 we see that, apart from multiplicative constants, the algebraic expression $\mathfrak{I}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \rightarrow \mathbf{R}$ of an unnormalized fundamental solution for the polyharmonic equation on $\mathbf{R}^d$ for d even, $k \geq d/2$, is given by

$$\mathfrak{I}_k^d(\mathbf{x}, \mathbf{x}') := \|\mathbf{x} - \mathbf{x}'\|^{2k-d} (\log \|\mathbf{x} - \mathbf{x}'\| - \beta_{k-d/2,d}). \quad (9)$$

By expressing $\mathfrak{I}_k^d$ in a rotationally-invariant coordinate system (6) we obtain

$$\begin{aligned} \mathfrak{I}_k^d(\mathbf{x}, \mathbf{x}') &= (2RR')^p \left[\frac{1}{2} \log(2RR') - \beta_{p,d} \right] [\chi - \cos(\phi - \phi')]^p \\ &\quad + \frac{1}{2} (2RR')^p [\chi - \cos(\phi - \phi')]^p \log [\chi - \cos(\phi - \phi')], \end{aligned} \quad (10)$$

where $p = k - d/2 \in \mathbf{N}_0$. For the polyharmonic equation on even-dimensional Euclidean space $\mathbf{R}^d$ with $1 \leq k \leq d/2 - 1$, apart from multiplicative constants, the algebraic expression for an unnormalized fundamental solution of the polyharmonic equation $\mathfrak{h}_k^d : (\mathbf{R}^d \times \mathbf{R}^d) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{R}^d\} \rightarrow \mathbf{R}$ is given by

$$\mathfrak{h}_k^d(\mathbf{x}, \mathbf{x}') := \|\mathbf{x} - \mathbf{x}'\|^{2k-d}.$$

By expressing $\mathfrak{h}_k^d$ in a rotationally-invariant coordinate system we obtain

$$\mathfrak{h}_k^d(\mathbf{x}, \mathbf{x}') = (2RR')^{-q} [\chi - \cos(\phi - \phi')]^{-q}, \quad (11)$$

where $q = 2k - d \in \mathbf{N}$.

By examining (10), (11), we see that for computation of Fourier expansions about the azimuthal separation angle $(\phi - \phi')$ of $\mathfrak{l}_k^d$ and $\mathfrak{h}_k^d$, all that is required is to compute the Fourier cosine series for the following three functions $f_\chi, h_\chi : \mathbf{R} \rightarrow (0, \infty)$ and $g_\chi : \mathbf{R} \rightarrow \mathbf{R}$ defined as

$$\begin{aligned} f_\chi(\psi) &:= (\chi - \cos \psi)^p, \\ g_\chi(\psi) &:= (\chi - \cos \psi)^p \log(\chi - \cos \psi), \\ h_\chi(\psi) &:= (\chi - \cos \psi)^{-q}, \end{aligned}$$

where $p \in \mathbf{N}_0$, $q \in \mathbf{N}$ and $\chi > 1$ is a fixed parameter. In a polyspherical coordinate system, the separation angle $\gamma \in [0, \pi]$, is the smallest angle measured between two arbitrary non-zero length position vectors $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$. It is defined through the relation

$$\cos \gamma := \frac{(\mathbf{x}, \mathbf{x}')}{\|\mathbf{x}\| \|\mathbf{x}'\|}, \quad (12)$$

using the Euclidean inner product and norm (cf. (5)).

The Fourier series of f_χ is given in Cohl & Dominici (2010) [5] (cf. (4.4) therein), namely for $p \in \mathbf{N}_0$,

$$(z - x)^p = (z^2 - 1)^{p/2} \sum_{n=0}^p \frac{\epsilon_n(-p)_n(p-n)!}{(p+n)!} P_p^n \left(\frac{z}{\sqrt{z^2 - 1}} \right) T_n(x), \quad (13)$$

where $\epsilon_n \in \{1, 2\}$ is the Neumann factor defined by $\epsilon_n := 2 - \delta_{n,0}$, $\delta_{n,0} \in \{0, 1\}$ is the Kronecker delta, and the Pochhammer symbol (rising factorial) $(\cdot)_n : \mathbf{C} \rightarrow \mathbf{C}$ for $n \in \mathbf{N}_0$ is defined by

$$(z)_n := \prod_{i=1}^n (z + i - 1).$$

We have used Whipple's formula in (13) (see for instance Olver *et al.* (2010) [21, (14.9.16), (14.9.17)]) to convert the associated Legendre function of the second kind Q_ν^μ appearing in [5] to the associated Legendre function of the first kind P_ν^μ . The associated Legendre function of the second kind $Q_\nu^\mu : \mathbf{C} \setminus (-\infty, 1] \rightarrow \mathbf{C}$, $\nu + \mu \notin -\mathbf{N}$, can be defined in terms of the Gauss hypergeometric function as follows (Olver *et al.* (2010) [21, (14.3.7) and section 14.21])

$$Q_\nu^\mu(z) := \frac{\sqrt{\pi} e^{i\pi\mu} \Gamma(\nu + \mu + 1) (z^2 - 1)^{\mu/2}}{2^{\nu+1} \Gamma(\nu + \frac{3}{2}) z^{\nu+\mu+1}} {}_2F_1 \left(\begin{matrix} \frac{\nu+\mu+2}{2}, \frac{\nu+\mu+1}{2} \\ \nu + \frac{3}{2} \end{matrix} ; \frac{1}{z^2} \right),$$

for $|z| > 1$ and by analytic continuation of the Gauss hypergeometric function elsewhere. Similarly, the associated Legendre function of the first kind can be defined using the Gauss hypergeometric function ((14.3.6) and §14.21(i) in Olver *et al.* (2010) [21])

$$P_\nu^\mu(z) := \frac{1}{\Gamma(1 - \mu)} \left(\frac{z+1}{z-1} \right)^{\mu/2} {}_2F_1 \left(\begin{matrix} -\nu, \nu+1 \\ 1 - \mu \end{matrix} ; \frac{1-z}{2} \right), \quad (14)$$

where $|1 - z| < 2$, and elsewhere by analytic continuation. The Gauss hypergeometric function ${}_2F_1 : \mathbf{C} \times \mathbf{C} \times (\mathbf{C} \setminus -\mathbf{N}_0) \times \{z \in \mathbf{C} : |z| < 1\} \rightarrow \mathbf{C}$ can be defined in terms of the following infinite series

$${}_2F_1 \left(\begin{matrix} a, b \\ c \end{matrix}; z \right) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{z^n}{n!} \quad (15)$$

(see for instance Chapter 15 in Olver *et al.* (2010) [21]), and elsewhere by analytic continuation.

The Fourier series of h_χ is given in Cohl & Dominici (2010) [5, (4.5)] (we have used the Whipple formula), namely for $p \in \mathbf{N}$,

$$\frac{1}{(z - x)^p} = \frac{(z^2 - 1)^{-p/2}}{(p - 1)!} \sum_{n=0}^{\infty} \epsilon_n (n + p - 1)! P_{p-1}^{-n} \left(\frac{z}{\sqrt{z^2 - 1}} \right) T_n(x). \quad (16)$$

In order to compute Fourier expansion of $\mathfrak{l}_k^d$ (10) in separable rotationally-invariant coordinate systems, all that remains is to determine the Fourier series of g_χ (see Cohl (2012) [4]). We will leave the discussion of Gauss hypergeometric orthogonal polynomial expansions for logarithmic fundamental solutions of the polyharmonic equation on $\mathbf{R}^d$ (from g_χ) to a later paper and from here on out we will only discuss expansions of power-law fundamental solutions (from f_χ, h_χ) of the polyharmonic equation.

3. Vilenkin's polyspherical coordinate systems on $\mathbf{R}^d$

In this section we review results in the literature concerning Vilenkin's polyspherical coordinates, which are subgroup type coordinates. Subgroup type coordinate systems on $\mathbf{R}^d$ are coordinate systems which can be described by a maximal chain of subgroups of the Euclidean group $E(d)$.

Polyspherical coordinates are specific subgroup type coordinate systems which are described by a radial coordinate $r \in [0, \infty)$ plus $(d - 1)$ -angles which together parametrize points on a $(d - 1)$ -dimensional hypersphere. First we will discuss a general procedure for constructing polyspherical coordinate systems called the “method of trees,” and then give some examples of polyspherical coordinates which will be used in the rest of the paper.

3.1. Method of trees

Subgroup type coordinates are coordinates in which the very useful method of trees (a graphical method for generating coordinate transformations for subgroup type coordinates) can be adopted. Corresponding to each of these subgroup chains are a set of tree diagrams. Each of these trees corresponds to a particular separable coordinate system in d -dimensions. See Izmet'sev *et al.* (1999) [15] (and below) for explicit parametrizations for the corresponding coordinate systems and trees. Describing polyspherical coordinate systems in terms of rooted trees was originally developed by

Vilenkin, Kuznetsov & Smorodinskiĭ (1965) [26] (see also Vilenkin (1968) [25]) and has since been used extensively by others in a variety of contexts (see for instance Izmet's'ev *et al.* (1999) [15], Izmet's'ev *et al.* (2001) [16], Kil'dyushov (1972) [18]).

In these rooted trees, there are two types of nodes, leaf nodes and branching nodes. For a coordinate system on $\mathbf{R}^d$, there are d leaf nodes, each corresponding to the particular Cartesian component of an arbitrary position vector $\mathbf{x} \in \mathbf{R}^d$. The branching nodes split into two separate branches, one up to the left and one up to the right. Each branch emanating from a branching node will end on either a leaf node or on another branching node. There are four possibilities for branching nodes (see figure 1).




Figure 1. This figure shows the possibilities from left to right for branching nodes of type **a**, **b**, **b'** and **c**. For type **a**, both branches end on a leaf node. The angle corresponding to this type of branching node is $\phi_a \in [0, 2\pi)$. For type **b**, the left branch ends on a leaf node and the right branch ends on a branching node. The angle corresponding to this type of branching node is $\theta_b \in [0, \pi]$. For type **b'**, the left-branch ends on a branching node and the right branch ends on a leaf node. The angle corresponding to this type of branching node is $\theta_{b'} \in [-\pi/2, \pi/2]$. For type **c**, both the left and right branches ends on branching nodes (cells of type **c** are only possible for $d \geq 4$). The angle corresponding to this type of branching node is $\vartheta_c \in [0, \pi/2]$.

A linear partial differential equation on $\mathbf{R}^d$ may admit solutions via the separation of variables method. If the method works, then it produces d -ordinary differential equations with $(d - 1)$ -separation constants which are determined by the conditions imposed on the problem being solved. The sets on which the separation constants are contained depend on the geometric structure of the problem being solved. For instance, separation of variables in polyspherical coordinates with $(d - 1)$ -angles, for Laplace's equation on $\mathbf{R}^d$ produces $(d - 1)$ -separation constants, each of which are called quantum numbers. The quantum numbers corresponding to these angles are all integers. This is due to the fact that a hypersphere is a compact manifold. If the separation constants are integers, then they are referred to as quantum numbers. Quantum numbers for a particular tree label the basis of separable solutions for Laplace's equation in that particular coordinate system. With each branching node of the tree, we associate a quantum number as well as an angle.

The quantum number corresponding to a (2π) -periodic (azimuthal) angle is called an azimuthal quantum number. Each azimuthal angle corresponds to a branching node of type **a**, and an azimuthal quantum number $m \in \mathbf{Z}$. A natural consequence of the

method of trees is that there must exist at least one azimuthal angle for each tree, and therefore also for each polyspherical coordinate system.

Branching nodes of type **b**, **b'** and **c**, as described above, are associated with angles, which in turn are associated with quantum numbers which we refer to as angular momentum quantum numbers $l \in \mathbf{N}_0$. The naming of these quantum numbers is due to the intimate connection between the properties of the rotation group $SO(d)$ and the quantum/classical description of angular momentum in physics. See for instance Chapter 10 in Fano & Rau (1996) [11].

There is always at least one branching node, the root, and all branching nodes correspond to a particular angle and quantum number. Let us associate each branching node with an angle and its corresponding quantum number. Parametrization of polyspherical coordinate systems are accomplished as follows. Starting at the root, traverse the tree upward until you reach the leaf node corresponding to x_i . The parametrization for x_i is given by the hyperspherical radius r multiplied by cosine or sine of each angle encountered as you traverse the tree upward until you reach the leaf node corresponding to x_i . If you branch upwards to the left or upwards to the right at each branching node, multiply by the cosine or sine of the corresponding angle respectively. This procedure produces the appropriate transformation from polyspherical coordinates to Cartesian coordinates.

There are large numbers of equivalent trees and an even larger number of possible trees, each with their own specific polyspherical coordinate system. The enumeration of these trees are characterized as follows. For $d, b_d \in \mathbf{N}$, let b_d be the total number of trees (polyspherical coordinate systems) on $\mathbf{R}^d$. Then $b_1 = 1$ is the number of possible 1-branch trees. A 1-branch tree corresponding to a polyspherical coordinate system does not exist in isolation. However, the concept of a 1-branch tree is useful. For instance, one can generate a new tree from a pre-existing tree by adding a single branch to either the left or to the right of a pre-existing tree and in order to perform the correct enumeration, one must incorporate 1-branch trees. The following recurrence relation gives the total number of possible polyspherical coordinate systems (trees) for arbitrary dimension

$$b_d = \sum_{i=1}^{d-1} b_i b_{d-i}. \quad (17)$$

Using the recurrence relation given by (17), the first few elements of the sequence are given by

$$(b_d : d \in \{2, \dots, 13\}) = (1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, 208012).$$

These numbers, or the way to count these types of trees, are given in terms of the Catalan numbers C_n (see for instance Sloane Integer Sequence A000108 [23] or p. 200 in Stanley (1999) [24]), i.e.,

$$b_d = C_{d-1} = \frac{1}{d} \binom{2d-2}{d-1},$$

where

$$\binom{n}{k} = \frac{n!}{k!(n-k)!}$$

is the binomial coefficient for $k, n \in \mathbf{N}_0$ with $0 \leq k \leq n$.

If $a_d \in \mathbf{N}$ is the total number of equivalence classes for equivalent trees for each dimension d , then the following recurrence relation gives the number of equivalence classes for arbitrary d

$$a_d = \begin{cases} \sum_{i=1}^{\lfloor d/2 \rfloor} a_i a_{d-i}, & \text{if } d \text{ odd} \\ \sum_{i=1}^{d/2-1} a_i a_{d-i} + \frac{1}{2} a_{d/2} (a_{d/2} + 1), & \text{if } d \text{ even.} \end{cases} \quad (18)$$

Using the recurrence relation given by (18), the first few elements of the sequence are given by

$$(a_d : d \in \{2, \dots, 13\}) = (1, 1, 2, 3, 6, 11, 23, 46, 98, 207, 451, 983).$$

These numbers, or the way to count these types of trees, are given in terms of the Wedderburn-Etherington numbers (see for instance Sloane Integer Sequence A001190 [23]). The equivalence classes are established by a left-right symmetry in the topology of the trees. The types of polyspherical coordinates that one can construct, for a given dimension, are described in Izmet'sev *et al.* (1999, 2001) [15, 16].

We use a left-to-right recursive naming language for our trees based on a depth-first search (see pp. 540-549 in Cormen *et al.* (2001) [9]). This naming language is given by listing the types of branching nodes available in a particular tree.

3.2. Vilenkin's polyspherical coordinate systems

A particular polyspherical coordinate system partitions $\mathbf{R}^d$ into a family of concentric $(d-1)$ -dimensional hyperspheres, each with a radius $r \in (0, \infty)$, on which all polyspherical coordinate systems for $\mathbf{S}_r^{d-1}$ may be used. One must also consider the limiting case for $r = 0$ to fill out all of $\mathbf{R}^d$. The hyperspherical radius is computed using Cartesian coordinates through

$$r^2 := \sum_{i=1}^d x_i^2.$$

A polyspherical coordinate system yields solutions to Laplace's equation on $\mathbf{R}^d$ through the separation of variables method.

The method of trees can be used to construct the cosine of the separation angle (12) in a direct manner. The cosine of the separation angle will be given by the sum of d -terms, each corresponding a leaf node of the tree. There is a unique path starting

from the root to each leaf node. The cosine of the separation angle can be constructed using the following formula

$$\cos \gamma = \sum_{i=1}^d \prod_{j=1}^{N_i} A_{i,j}(\psi_{i,j}) A_{i,j}(\psi'_{i,j}),$$

where N_i is the number of branching nodes encountered from the root to the leaf node, $A_{i,j} : \mathbf{R} \rightarrow [-1, 1]$ is either the trigonometric cosine or sine function depending respectively on whether the left branch or right branch is chosen, and $\psi_{i,j} \in \mathbf{R}$ is the angle corresponding to the j th branching node for each i th leaf node. The formula for the cosine of the separation angle is unique for each tree.

3.3. Low dimensional examples of Vilenkin's polyspherical coordinate systems

The simplest example of a polyspherical coordinate system on $\mathbf{R}^d$ occurs for $d = 2$ (polar coordinates) where there is one branching node (the root) and two leaf nodes (see figures 2a and 5a). The left-branch ends on the leaf node corresponding to x_1 and the right branch ends on the leaf node corresponding to x_2 . Therefore for $d = 2$ we have the following coordinate system whose transformation formulae to Cartesian coordinates is given by

$$\left. \begin{aligned} x_1 &= r \cos \phi \\ x_2 &= r \sin \phi \end{aligned} \right\}, \quad (19)$$

where $r \in [0, \infty)$ and $\phi \in [0, 2\pi)$. This coordinate system is the only member of its equivalence class (cf. (18)). Using our naming procedure based on a depth-first search, we refer to this tree as type **a**. These coordinates are adapted to the canonical subgroup chain $E(2) \supset O(2)$. The cosine of the separation angle is given by

$$\cos \gamma = \cos(\phi - \phi'), \quad (20)$$

and corresponding to the angle ϕ is the quantum number $m \in \mathbf{Z}$ (see figure 2a).




Figure 2. Tree diagrams for two and three dimensional polyspherical coordinate systems of type: (a) **a**, (b) **ba**, (c) **b'a**.

In $d = 3$ there are 2 possible topological trees, each corresponding to one of two different trees. Both coordinate systems are adapted to the canonical subgroup

chain $E(3) \supset O(3) \supset O(2)$. The first tree (see figure 2b) corresponds to the following coordinate system whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta \\ x_2 &= r \sin \theta \cos \phi \\ x_3 &= r \sin \theta \sin \phi \end{aligned} \right\}, \quad (21)$$

where $\theta \in [0, \pi]$, $\phi \in [0, 2\pi)$. We call this polyspherical coordinate system *standard spherical coordinates*. Using our naming procedure based on a depth-first search, we refer to this tree as type **ba**. The cosine of the separation angle is given by

$$\cos \gamma = \cos \theta \cos \theta' + \sin \theta \sin \theta' \cos(\phi - \phi'), \quad (22)$$

and corresponding to the angles $\theta \in [0, \pi]$ and $\phi \in [0, 2\pi)$ are the quantum numbers $l \in \mathbf{N}_0$ and $m \in \mathbf{Z}$ respectively.

The second tree (see figure 2c) corresponds to the following coordinate system whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta \cos \phi \\ x_2 &= r \cos \theta \sin \phi \\ x_3 &= r \sin \theta \end{aligned} \right\}, \quad (23)$$

where $\theta \in [-\pi/2, \pi/2]$, $\phi \in [0, 2\pi)$. This tree is of type **b'a**. The cosine of the separation angle is given by

$$\cos \gamma = \sin \theta \sin \theta' + \cos \theta \cos \theta' \cos(\phi - \phi'), \quad (24)$$

and corresponding to the angles θ and ϕ are the quantum numbers $l \in \mathbf{N}_0$ and $m \in \mathbf{Z}$ respectively. The coordinate systems of type **ba** and **b'a** belong to the same equivalence class.

In $d = 4$ there are five possible topological trees for polyspherical coordinates, each corresponding to a different tree. The first four coordinate systems are adapted to the canonical subgroup chain $E(4) \supset O(4) \supset O(3) \supset O(2)$. The first tree (see figures 3a and 4) corresponds to the following coordinate system, whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta_1 \\ x_2 &= r \sin \theta_1 \cos \theta_2 \\ x_3 &= r \sin \theta_1 \sin \theta_2 \cos \phi \\ x_4 &= r \sin \theta_1 \sin \theta_2 \sin \phi \end{aligned} \right\}, \quad (25)$$

where $\theta_1, \theta_2 \in [0, \pi]$. Using our naming procedure based on a depth-first search, this tree is of type **b²a**. The cosine of the separation angle is given by

$$\cos \gamma = \cos \theta_1 \cos \theta'_1 + \sin \theta_1 \sin \theta'_1 (\cos \theta_2 \cos \theta'_2 + \sin \theta_2 \sin \theta'_2 \cos(\phi - \phi')). \quad (26)$$




Figure 3. Tree diagrams for four dimensional polyspherical coordinate systems of type: (a) $\mathbf{b}^2\mathbf{a}$, (b) $\mathbf{bb}'\mathbf{a}$, (c) $\mathbf{b}'\mathbf{ba}$, (d) $\mathbf{b}'^2\mathbf{a}$, (e) $\mathbf{ca}^2$.

The second tree (see figure 3b) corresponds to the following coordinate system whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta_1 \\ x_2 &= r \sin \theta_1 \cos \theta_2 \cos \phi \\ x_3 &= r \sin \theta_1 \cos \theta_2 \sin \phi \\ x_4 &= r \sin \theta_1 \sin \theta_2 \end{aligned} \right\}, \quad (27)$$

where $\theta_1 \in [0, \pi]$ and $\theta_2 \in [-\pi/2, \pi/2]$. This tree is of type $\mathbf{bb}'\mathbf{a}$. The cosine of the separation angle is given by

$$\cos \gamma = \cos \theta_1 \cos \theta'_1 + \sin \theta_1 \sin \theta'_1 (\sin \theta_2 \sin \theta'_2 + \cos \theta_2 \cos \theta'_2 \cos(\phi - \phi')). \quad (28)$$

The third tree (see figure 3c) corresponds to the following coordinate system, whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta_1 \cos \theta_2 \\ x_2 &= r \cos \theta_1 \sin \theta_2 \cos \phi \\ x_3 &= r \cos \theta_1 \sin \theta_2 \sin \phi \\ x_4 &= r \sin \theta_1 \end{aligned} \right\}, \quad (29)$$

where $\theta_1 \in [-\pi/2, \pi/2]$ and $\theta_2 \in [0, \pi]$. This tree is of type $\mathbf{b}'\mathbf{ba}$. The cosine of the separation angle is given by

$$\cos \gamma = \sin \theta_1 \sin \theta'_1 + \cos \theta_1 \cos \theta'_1 (\cos \theta_2 \cos \theta'_2 + \sin \theta_2 \sin \theta'_2 \cos(\phi - \phi')). \quad (30)$$

The fourth tree (see figure 3d) corresponds to the following coordinate system, whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta_1 \cos \theta_2 \cos \phi \\ x_2 &= r \cos \theta_1 \cos \theta_2 \sin \phi \\ x_3 &= r \cos \theta_1 \sin \theta_2 \\ x_4 &= r \sin \theta_2 \end{aligned} \right\}, \quad (31)$$

where $\theta_1, \theta_2 \in [-\pi/2, \pi/2]$. This tree is of type $\mathbf{b}'^2\mathbf{a}$. The cosine of the separation angle is given by

$$\cos \gamma = \sin \theta_1 \sin \theta'_1 + \cos \theta_1 \cos \theta'_1 (\sin \theta_2 \sin \theta'_2 + \cos \theta_2 \cos \theta'_2 \cos(\phi - \phi')). \quad (32)$$

The fifth tree (see figures 3e and 5b) corresponds to the following coordinate system (Hopf coordinates) whose transformation formulae to Cartesian coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \vartheta \cos \phi_1 \\ x_2 &= r \cos \vartheta \sin \phi_1 \\ x_3 &= r \sin \vartheta \cos \phi_2 \\ x_4 &= r \sin \vartheta \sin \phi_2 \end{aligned} \right\}, \quad (33)$$

where $\vartheta \in [0, \pi/2]$ and $\phi_1, \phi_2 \in [0, 2\pi)$. This tree is of type $\mathbf{ca}^2$. This coordinate system is adapted to the canonical subgroup chain $E(4) \supset O(4) \supset O(2) \otimes O(2)$. The cosine of the separation angle is given by

$$\cos \gamma = \cos \vartheta \cos \vartheta' \cos(\phi_1 - \phi'_1) + \sin \vartheta \sin \vartheta' \cos(\phi_2 - \phi'_2). \quad (34)$$

3.4. High dimensional examples of Vilenkin's polyspherical coordinate systems

There are many choices for polyspherical coordinates on $\mathbf{R}^d$, suitably defined for any number of dimensions $d \geq 2$. Note that these also give polyspherical coordinate systems on $\mathbf{S}^{d-1}$ when $r = 1$. The simplest example of a polyspherical coordinate system on $\mathbf{R}^d$, valid for high dimensions, is what we will refer to as *standard polyspherical coordinates* (see figure 4). Standard polyspherical coordinates are a generalization of (21, type $\mathbf{ba}$) and (25, type $\mathbf{b}^2\mathbf{a}$). This coordinate system (and all those which belong to its same equivalence class) is adapted to the canonical subgroup chain $E(d) \supset O(d) \supset O(d-1) \supset \cdots \supset O(2)$. The transformation formulae to Cartesian coordinates for standard polyspherical coordinates are given by

$$\left. \begin{aligned} x_1 &= r \cos \theta_1 \\ x_2 &= r \sin \theta_1 \cos \theta_2 \\ x_3 &= r \sin \theta_1 \sin \theta_2 \cos \theta_3 \\ &\vdots \\ x_{d-2} &= r \sin \theta_1 \cdots \sin \theta_{d-3} \cos \theta_{d-2} \\ x_{d-1} &= r \sin \theta_1 \cdots \sin \theta_{d-3} \sin \theta_{d-2} \cos \phi \\ x_d &= r \sin \theta_1 \cdots \sin \theta_{d-3} \sin \theta_{d-2} \sin \phi \end{aligned} \right\}, \quad (35)$$

where $\theta_i \in [0, \pi]$ for $i \in \{1, \dots, d-2\}$ and $\phi \in [0, 2\pi)$. Using our naming procedure based on a depth-first search, this tree is of type $\mathbf{b}^{d-2}\mathbf{a}$. These are the standard polyspherical coordinates that are usually adopted in multi-variable calculus and in physical applications (see for instance Fano & Rau (1996) [11]). In these coordinates the cosine of the separation angle is given by

$$\cos \gamma = \sum_{i=1}^{d-2} \cos \theta_i \cos \theta'_i \prod_{j=1}^{i-1} \sin \theta_j \sin \theta'_j + \cos(\phi - \phi') \prod_{i=1}^{d-2} \sin \theta_i \sin \theta'_i.$$




Figure 4. Tree diagram for d -dimensional standard polyspherical coordinates of type $\mathbf{b}^{d-2}\mathbf{a}$ (standard polyspherical coordinates).

Another example of a polyspherical coordinate system which is valid for large dimensions is what we will refer to as *generalized Hopf coordinates* (see figure 5). These coordinates, valid on $\mathbf{R}^{2^q}$ for $q \geq 1$, generalize two-dimensional polar coordinates (19, type $\mathbf{a}$) and four-dimensional Hopf coordinates (33, type $\mathbf{ca}^2$). These coordinates are unique in that they correspond to the only trees which contain only themselves in their equivalence class (see (18)). These coordinate systems have separated harmonic eigenfunctions which are given in terms of complex exponentials, and for $q \geq 2$, non-symmetric Jacobi polynomials (see figure 5). These coordinates are adapted to the canonical subgroup chain

$$\begin{aligned} E(d) &\supset O(d) \supset O(d-1) \otimes O(d-1) \\ &\supset O(d-2) \otimes O(d-2) \otimes O(d-2) \otimes O(d-2) \supset \dots \supset O(2) \otimes \dots \otimes O(2). \end{aligned}$$




Figure 5. This figure is a tree diagram for polyspherical generalized Hopf coordinates of type $\mathbf{V}_{2^q}$ on $\mathbf{R}^{2^q}$ with $q = 1, 2, 3, 4$ for (a), (b), (c), and (d) respectively. The first $(2^{q-1} - 1)$ -branching nodes are of type **c** which correspond to the angles $\vartheta_i \in [0, \frac{\pi}{2}]$ and quantum numbers $l_i \in \mathbf{N}_0$. The following (2^q) -branching nodes are of type **a** which correspond to the angles $\phi_i \in [0, 2\pi)$ and quantum numbers $m_i \in \mathbf{Z}$. These coordinates correspond to Transformation (36).

This coordinate system is suitably defined for dimensions $d = 2^q$ for $q \geq 1$. The transformation formulae to Cartesian coordinates are given by

$$\begin{aligned}
 x_1 &= r \cos \vartheta_1 \cos \vartheta_2 \cos \vartheta_4 \cos \vartheta_8 \cdots \cos \vartheta_{2^{q-2}} \cos \phi_1 \\
 x_2 &= r \cos \vartheta_1 \cos \vartheta_2 \cos \vartheta_4 \cos \vartheta_8 \cdots \cos \vartheta_{2^{q-2}} \sin \phi_1 \\
 &\vdots \\
 x_{2^{q-1}-1} &= r \cos \vartheta_1 \sin \vartheta_2 \sin \vartheta_5 \sin \vartheta_{11} \cdots \sin \vartheta_{3 \cdot 2^{q-3}-1} \cos \phi_{2^{q-2}} \\
 x_{2^{q-1}} &= r \cos \vartheta_1 \sin \vartheta_2 \sin \vartheta_5 \sin \vartheta_{11} \cdots \sin \vartheta_{3 \cdot 2^{q-3}-1} \sin \phi_{2^{q-2}} \\
 x_{2^{q-1}+1} &= r \sin \vartheta_1 \cos \vartheta_3 \cos \vartheta_6 \cos \vartheta_{12} \cdots \cos \vartheta_{3 \cdot 2^{q-3}} \cos \phi_{2^{q-2}+1} \\
 x_{2^{q-1}+2} &= r \sin \vartheta_1 \cos \vartheta_3 \cos \vartheta_6 \cos \vartheta_{12} \cdots \cos \vartheta_{3 \cdot 2^{q-3}} \sin \phi_{2^{q-2}+1} \\
 &\vdots \\
 x_{2^q-1} &= r \sin \vartheta_1 \sin \vartheta_3 \sin \vartheta_7 \sin \vartheta_{17} \cdots \sin \vartheta_{3 \cdot 2^{q-1}-1} \cos \phi_{2^{q-1}}
 \end{aligned}
 \quad \left. \vphantom{\begin{aligned} x_1 \\ x_2 \\ \vdots \\ x_{2^{q-1}-1} \\ x_{2^{q-1}} \\ x_{2^{q-1}+1} \\ x_{2^{q-1}+2} \\ \vdots \\ x_{2^q-1} \end{aligned}} \right\}, \quad (36)$$

where $\vartheta_i \in [0, \frac{\pi}{2}]$ for $i = 1, \dots, 2^{q-1} - 1$ and $\phi_i \in [0, 2\pi)$ for $i = 1, \dots, 2^{q-1}$. Using our naming procedure based on a depth-first search (see section 3.1), these coordinates are of type

$$\mathbf{V}_{2^q} = \mathbf{c} \mathbf{V}_{2^{q-1}} \mathbf{V}_{2^{q-1}},$$

where $q \in \mathbf{N}$, with type $\mathbf{V}_2 = \mathbf{a}$ (polar) coordinates on $\mathbf{R}^2$ (see figures 2a and 5a). The first non-trivial example of generalized Hopf coordinates is $\mathbf{V}_4 = \mathbf{c} \mathbf{V}_2 \mathbf{V}_2 = \mathbf{c} \mathbf{a}^2$ (Hopf coordinates, see figures 3e and 5b). Other examples of generalized Hopf coordinates are of type $\mathbf{V}_8 = \mathbf{c} \mathbf{V}_4 \mathbf{V}_4 = \mathbf{c} \mathbf{c} \mathbf{a}^2 \mathbf{c} \mathbf{a}^2$ and of type $\mathbf{V}_{16} = \mathbf{c} \mathbf{V}_8 \mathbf{V}_8 = \mathbf{c} \mathbf{c} \mathbf{c} \mathbf{a}^2 \mathbf{c} \mathbf{a}^2 \mathbf{c} \mathbf{c} \mathbf{a}^2 \mathbf{c} \mathbf{a}^2$ coordinates (for corresponding tree diagrams, see figures 5c and 5d).

The cosine of the separation angle in this coordinate system may be given as follows. Consider the symbol ${}_q \mathbf{G}_s^r \in [-1, 1]$ with $0 \leq s \leq q$ and $1 \leq r \leq 2^q - 1$. If this symbol is given recursively by

$${}_q \mathbf{G}_s^r = \cos \vartheta_{r-1+2^{q-s}} \cos \vartheta'_{r-1+2^{q-s}} {}_q \mathbf{G}_{s-1}^{2r-1} + \sin \vartheta_{r-1+2^{q-s}} \sin \vartheta'_{r-1+2^{q-s}} {}_q \mathbf{G}_{s-1}^{2r},$$

with ${}_q \mathbf{G}_0^i = 1$, then the cosine of the separation angle is given by

$$\cos \gamma = {}_q \mathbf{G}_q^1$$

(cf. (20), (34)). Note that through the identification $\phi_i = \vartheta_{i-1+2^{q-1}}$ with $i = 1, \dots, 2^{q-1}$, then ${}_q \mathbf{G}_1^i = \cos(\phi_i - \phi'_i)$, and using this one may stop this recursion at $s = 1$.

4. Hyperspherical harmonics in polyspherical coordinates

The general basis functions that one gets by putting coordinates on the d -dimensional hypersphere $\mathbf{S}^d$, can be specified as solutions to the angular part of Laplace's equation on $\mathbf{R}^{d+1}$. These correspond to separated solutions of Laplace's equation, using the Laplace-Beltrami operator on the hypersphere $\mathbf{S}^d$. The following numbers are associated with each cell $m \in \mathbf{Z}$, $l, l_\alpha, l_\beta \in \mathbf{N}_0$ (see figure 1 in section 3.2). The number of vertices above each branching node l_α and l_β are represented by S_α and S_β respectively. The numbers m, l, l_α, l_β label representations for the corresponding rotation subgroup in the subgroup chain, i.e., angular momentum type quantum numbers (see Izmet'ev *et al.* (2001) [16]).

The following eigenfunctions are generated at each branching node for normalized hyperspherical harmonics in polyspherical coordinates using the method of trees (please refer to figure 1):

- Type **a**:

$$\Psi_m(\phi_a) = \frac{1}{\sqrt{2\pi}} e^{im\phi_a}; \quad m \in \mathbf{Z}, \quad \phi \in [0, 2\pi). \quad (37)$$

- Type **b**:

$$\left. \begin{aligned} \Psi_{n,l_\beta}^\alpha(\theta_b) &= N_n^{\alpha,\alpha} (\sin \theta_b)^{l_\beta} P_n^{(\alpha,\alpha)}(\cos \theta_b); \\ n &= l - l_\beta, \quad \alpha = l_\beta + \frac{S_\beta}{2}, \quad n \in \mathbf{N}_0, \quad \theta_b \in [0, \pi]. \end{aligned} \right\} \quad (38)$$

- Type **b'**:

$$\left. \begin{aligned} \Psi_{n,l_\alpha}^\beta(\theta_{b'}) &= N_n^{\beta,\beta}(\cos \theta_{b'})^{l_\alpha} P_n^{(\beta,\beta)}(\sin \theta_{b'}); \\ n = l - l_\alpha, \quad \beta &= l_\alpha + \frac{S_\alpha}{2}, \quad n \in \mathbf{N}_0, \quad \theta_{b'} \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]. \end{aligned} \right\} \quad (39)$$

- Type **c**:

$$\left. \begin{aligned} \Psi_{n,l_\alpha,l_\beta}^{\alpha,\beta}(\vartheta_c) &= 2^{(\alpha+\beta)/2+1} N_n^{\alpha,\beta}(\sin \vartheta_c)^{l_\beta} (\cos \vartheta_c)^{l_\alpha} P_n^{(\beta,\alpha)}(\cos 2\vartheta_c); \\ n = \frac{1}{2}(l - l_\alpha - l_\beta), \quad \alpha &= l_\alpha + \frac{S_\alpha}{2}, \quad \beta = l_\beta + \frac{S_\beta}{2}, \quad n \in \mathbf{N}_0, \quad \vartheta_c \in \left[0, \frac{\pi}{2}\right]. \end{aligned} \right\} \quad (40)$$

We refer to the quantum number $n \in \mathbf{N}_0$ in (40) as the *surrogate* quantum number to $l \in \mathbf{N}_0$. The Jacobi polynomials $P_n^{(\alpha,\beta)} : \mathbf{C} \rightarrow \mathbf{C}$ can be defined in terms of the terminating Gauss hypergeometric series as follows (Olver *et al.* (2010) [21, (18.5.7)])

$$P_n^{(\alpha,\beta)}(z) := \frac{(\alpha+1)_n}{n!} {}_2F_1 \left(\begin{matrix} -n, n+\alpha+\beta+1 \\ \alpha+1 \end{matrix}; \frac{1-z}{2} \right),$$

for $n \in \mathbf{N}_0$, and $\alpha, \beta > -1$. The normalization constants $N_n^{\alpha,\beta} \in (0, \infty)$ are given by

$$N_n^{\alpha,\beta} = \sqrt{\frac{(2n+\alpha+\beta+1)\Gamma(n+\alpha+\beta+1)n!}{2^{\alpha+\beta+1}\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}},$$

such that

$$\int_{-1}^1 P_m^{(\alpha,\beta)}(x) P_n^{(\alpha,\beta)}(x) (1-x)^\alpha (1+x)^\beta dx = \frac{\delta_{m,n}}{(N_n^{\alpha,\beta})^2},$$

where $\alpha, \beta > -1$ (see (7.391.1) in Gradshteyn & Ryzhik (2007) [13]).

Notice that the eigenfunctions for cells of type **b** and **b'** can be expressed in terms of Gegenbauer polynomials using (see (6.4.9) in Andrews, Askey & Roy (1999) [2])

$$C_n^\nu(x) = \frac{(2\nu)_n}{(\nu + \frac{1}{2})_n} P_n^{(\nu-1/2, \nu-1/2)}(x), \quad (41)$$

where $\nu > -1/2$. Therefore we can re-write (38) as

$$\Psi_{n,l_\beta}^\alpha(\theta_b) = \frac{(2\alpha)!}{\Gamma(\alpha+1)} \sqrt{\frac{(2\alpha+2n+1)n!}{2^{2\alpha+1}(2\alpha+n)!}} (\sin \theta_b)^{l_\beta} C_n^{\alpha+1/2}(\cos \theta_b), \quad (42)$$

and (39) as

$$\Psi_{n,l_\alpha}^\beta(\theta_{b'}) = \frac{(2\beta)!}{\Gamma(\beta+1)} \sqrt{\frac{(2\beta+2n+1)n!}{2^{2\beta+1}(2\beta+n)!}} (\sin \theta_{b'})^{l_\alpha} C_n^{\beta+1/2}(\cos \theta_{b'}). \quad (43)$$

Note that even though α, β are not necessarily integers, $2\alpha, 2\beta \in \mathbf{N}_0$.

4.1. Examples of hyperspherical harmonics in polyspherical coordinate systems

With the simplest example, polar coordinates (see figure 2a, Transformation (19)), the normalized harmonics are given by

$$Y_m(\phi) = \frac{e^{im\phi}}{\sqrt{2\pi}}.$$

In $d = 3$ there are two possible polyspherical coordinate systems, that of type **ba** (21) and **b'a** (23). In both cases there are two branching nodes. First we treat the type **ba** coordinate system (see figure 2b). Using (38) we see that $\alpha = m$, $n = l - m$, $l_\beta = m$, and $S_\beta = 0$ since there are no vertices above the branching node m . Through reduction and multiplication by the ϕ eigenfunction, the normalized spherical harmonics are

$$Y_{l,m}(\theta, \phi) = (-1)^m \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\cos \theta) e^{im\phi}, \quad (44)$$

where $P_\nu^\mu : (-1, 1) \rightarrow \mathbf{C}$ is the Ferrers function of the first kind (associated Legendre function of the first kind on the cut) through [21, (14.3.1)], defined as

$$P_\nu^\mu(x) := \frac{1}{\Gamma(1-\mu)} \left(\frac{1+x}{1-x} \right)^{\mu/2} {}_2F_1 \left(\begin{matrix} -\nu, \nu+1 \\ 1-\mu \end{matrix}; \frac{1-x}{2} \right). \quad (45)$$

Notice we have used (cf. (8.936.2) in Gradshteyn & Ryzhik (2007)) [13])

$$C_{l-m}^{m+1/2}(x) = \frac{(-1)^m (1-x^2)^{-m/2}}{(2m-1)!!} P_l^m(x), \quad (46)$$

with (42) to reduce the Gegenbauer polynomial to a Ferrers function of the first kind. The functions $Y_{l,m}$ are called *standard spherical harmonics*.

Now we treat the type **b'a** coordinate system (see figure 2c). Using (39) we see that $\alpha = m$, $n = l - m$, $l_\alpha = m$, and $S_\alpha = 0$ since there are no vertices above the branching node m . Through reduction and multiplication by the ϕ eigenfunction, the normalized spherical harmonics are

$$Y_{l,m}(\theta, \phi) = (-1)^m \sqrt{\frac{2l+1}{4\pi} \frac{(l-m)!}{(l+m)!}} P_l^m(\sin \theta) e^{im\phi}, \quad (47)$$

where we have used (43) and (46). The range of θ in (44) and (47) is taken to be $\theta \in [0, \pi]$ or $\theta \in [-\pi/2, \pi/2]$ (for nodes of type **b** and **b'**) respectively.

In $d = 4$ there are five possible polyspherical coordinate systems, those of type **b²a** (25), **bb'a** (27), **b'ba** (29), **b'²a** (31), and **ca²** (33). The normalized hyperspherical harmonics in type **b²a** coordinates (see figure 3a) are

$$Y_{l_1, l_2, m}(\theta_1, \theta_2, \phi) = \frac{(-1)^m (2l_2)!!}{\pi} \sqrt{\frac{(2l_2+1)(l_1+1)(l_1-l_2)!(l_2-m)!}{2(l_1+l_2+1)!(l_2+m)!}} \\ \times (\sin \theta_1)^{l_2} C_{l_1-l_2}^{l_2+1}(\cos \theta_1) P_{l_2}^m(\cos \theta_2) e^{im\phi}, \quad (48)$$

in type **bb'a** coordinates (see figure 3b)

$$Y_{l_1, l_2, m}(\theta_1, \theta_2, \phi) = \frac{(-1)^m (2l_2)!!}{\pi} \sqrt{\frac{(2l_2 + 1)(l_1 + 1)(l_1 - l_2)!(l_2 - m)!}{2(l_1 + l_2 + 1)!(l_2 + m)!}} \\ \times (\sin \theta_1)^{l_2} C_{l_1 - l_2}^{l_2 + 1}(\cos \theta_1) P_{l_2}^m(\sin \theta_2) e^{im\phi},$$

in type **b'ba** coordinates (see figure 3c)

$$Y_{l_1, l_2, m}(\theta_1, \theta_2, \phi) = \frac{(-1)^m (2l_2)!!}{\pi} \sqrt{\frac{(2l_2 + 1)(l_1 + 1)(l_1 - l_2)!(l_2 - m)!}{2(l_1 + l_2 + 1)!(l_2 + m)!}} \\ \times (\cos \theta_1)^{l_2} C_{l_1 - l_2}^{l_2 + 1}(\sin \theta_1) P_{l_2}^m(\cos \theta_2) e^{im\phi},$$

in type **b'²a** coordinates (see figure 3d)

$$Y_{l_1, l_2, m}(\theta_1, \theta_2, \phi) = \frac{(-1)^m (2l_2)!!}{\pi} \sqrt{\frac{(2l_2 + 1)(l_1 + 1)(l_1 - l_2)!(l_2 - m)!}{2(l_1 + l_2 + 1)!(l_2 + m)!}} \\ \times (\cos \theta_1)^{l_2} C_{l_1 - l_2}^{l_2 + 1}(\sin \theta_1) P_{l_2}^m(\sin \theta_2) e^{im\phi},$$

and in type **ca²** (see figure 3e)

$$Y_{l, m_1, m_2}(\vartheta, \phi_1, \phi_2) = \frac{e^{i(m_1 \phi_1 + m_2 \phi_2)}}{\pi} \sqrt{\frac{l + 1}{2} \frac{[\frac{1}{2}(l + |m_1| + |m_2|)]! [\frac{1}{2}(l - |m_1| - |m_2|)]!}{[\frac{1}{2}(l - |m_1| + |m_2|)]! [\frac{1}{2}(l + |m_1| - |m_2|)]!}} \\ \times (\sin \vartheta)^{|m_2|} (\cos \vartheta)^{|m_1|} P_{(l - |m_1| - |m_2|)/2}^{(|m_2|, |m_1|)}(\cos 2\vartheta), \quad (49)$$

with the restriction to the parameter space given by $\frac{1}{2}(l - |m_1| - |m_2|) \in \mathbf{N}_0$. Using the surrogate quantum number $n \in \mathbf{N}_0$, defined in terms of $l \in \mathbf{N}_0$ such that $2n = l - |m_1| - |m_2|$, we can more conveniently express the normalized hyperspherical harmonics in type **ca²** coordinates, namely

$$Y_{n, m_1, m_2}(\vartheta, \phi_1, \phi_2) = \frac{e^{i(m_1 \phi_1 + m_2 \phi_2)}}{\pi} \sqrt{\frac{(2n + |m_1| + |m_2| + 1)(n + |m_1| + |m_2|)! n!}{2(n + |m_1|)!(n + |m_2|)!}} \\ \times (\sin \vartheta)^{|m_2|} (\cos \vartheta)^{|m_1|} P_n^{(|m_2|, |m_1|)}(\cos 2\vartheta).$$

For arbitrary dimensions, we can use standard polyspherical coordinates (35) to construct normalized hyperspherical harmonics. The polyspherical harmonics corresponding to this coordinate system are basis functions for the irreducible representations of $O(d)$ (see Vilenkin (1968) [25]). In terms of these coordinates, the normalized hyperspherical harmonics are

$$Y_l^K(\hat{\mathbf{x}}) = \frac{e^{im\phi}}{\sqrt{2\pi}} \prod_{j=1}^{d-2} \Theta_j^d(l_j, l_{j+1}; \theta_j), \quad (50)$$

where $\hat{\mathbf{x}} \in \mathbf{S}^{d-1}$, $K = \{l_2, l_3, \dots, l_{d-1}\}$,

$$l = l_1 \geq l_2 \geq l_3 \geq \dots \geq l_{d-3} \geq l_{d-2} = \ell \geq l_{d-1} = |m| \geq 0, \quad (51)$$

and $\Theta_j^d : \mathbf{N}_0^2 \times [0, \pi] \rightarrow \mathbf{R}$ is defined by

$$\Theta_j^d(l_j, l_{j+1}; \theta_j) := \frac{\Gamma\left(l_{j+1} + \frac{d-j+1}{2}\right)}{2l_{j+1} + d - j - 1} \sqrt{\frac{2^{2l_{j+1}+d-j-1}(2l_j + d - j - 1)(l_j - l_{j+1})!}{\pi(l_j + l_{j+1} + d - j - 2)!}} \times (\sin \theta_j)^{l_{j+1}} C_{l_j - l_{j+1}}^{l_{j+1} + (d-j-1)/2}(\cos \theta_j). \quad (52)$$

The computation of (52) is a straightforward consequence of (38), and doing the proper node counting for S_β , in the tree depicted in figure 4. The normalized spherical harmonics (44) are the simplest example of these harmonics.

In generalized Hopf coordinates of type $\mathbf{V}_{2^p}$ coordinates with $p \geq 1$ (see figure 5), the normalized hyperspherical harmonics $Y_{l_1}^K : \mathbf{S}^{2^p-1} \rightarrow \mathbf{C}$ can be given more conveniently using the surrogate quantum numbers $n_q \in \mathbf{N}_0$ which are connected to $l_q \in \mathbf{N}_0$ through the relation $l_q = l_\alpha + l_\beta + 2n_q$ such that $1 \leq q \leq 2^{p-1} - 1$. The orthonormal polyspherical harmonics, which can be generated using the method of trees (see section 3.1), are

$$Y_{l_1}^K(\hat{\mathbf{x}}) = \frac{e^{i(m_1\phi_1 + \dots + m_{2^{p-1}}\phi_{2^{p-1}})}}{\sqrt{2\pi^{2^{p-2}}}} \prod_{j=1}^{p-1} \prod_{s=1}^{2^{j-1}} \Upsilon_q^p \left(\begin{matrix} n_q \\ l_\alpha, l_\beta \end{matrix}; \vartheta_q \right), \quad (53)$$

where $K = \{l_2, \dots, l_{2^{p-1}-1}, m_1, \dots, m_{2^{p-1}}\}$ and $\Upsilon_q^p : \mathbf{N}_0^3 \times [0, \frac{\pi}{2}] \rightarrow \mathbf{R}$ is defined by

$$\Upsilon_q^p \left(\begin{matrix} n_q \\ l_\alpha, l_\beta \end{matrix}; \vartheta_q \right) := \sqrt{\frac{(2n_q + \alpha + \beta + 1)(n_q + \alpha + \beta)! n_q!}{(n_q + \alpha)! (n_q + \beta)!}} \times (\cos \vartheta_q)^{l_\alpha} (\sin \vartheta_q)^{l_\beta} P_{n_q}^{(\beta, \alpha)}(\cos(2\vartheta_q)),$$

with $q = 2^{j-1} + s - 1$, $l_\alpha = l_{2q}$, $l_\beta = l_{2q+1}$, $\alpha = l_{2q} - 1 + 2^{p-2-\lfloor \log_2 q \rfloor}$, and $\beta = l_{2q+1} - 1 + 2^{p-2-\lfloor \log_2 q \rfloor}$. Note that we use the identification

$$(l_{2^{p-1}}, \dots, l_{2^{p-1}-1}) = (|m_1|, \dots, |m_{2^{p-1}}|).$$

5. Jacobi polynomial and limiting expansions for the Euler kernel

In this section we derive Jacobi, Gegenbauer and Chebyshev polynomial of the first kind series expansions of the Euler kernel $(z - x)^{-\nu}$. These series expansions are used to obtain azimuthal Fourier and hyperspherical harmonic expansions of a fundamental solution of the polyharmonic equation in Euclidean space.

Theorem 5.1. *Let $\alpha, \beta > -1$, $x, z, \nu \in \mathbf{C}$, with $z \in \mathbf{C} \setminus (-\infty, 1]$ on any ellipse with foci at ± 1 and x in the interior of that ellipse. Then*

$$\frac{1}{(z - x)^\nu} = \frac{(z - 1)^{\alpha+1-\nu} (z + 1)^{\beta+1-\nu}}{2^{\alpha+\beta+1-\nu}} \times \sum_{n=0}^{\infty} \frac{(\alpha + \beta + 2n + 1) \Gamma(\alpha + \beta + n + 1) (\nu)_n}{\Gamma(\alpha + 1 + n) \Gamma(\beta + 1 + n)} Q_{n+\nu-1}^{(\alpha+1-\nu, \beta+1-\nu)}(z) P_n^{(\alpha, \beta)}(x). \quad (54)$$

If one substitutes $z = (1 + \rho^2)/(2\rho)$ in (54) with $0 < |\rho| < 1$, then one obtains an alternate expression

$$\frac{1}{(1 + \rho^2 - 2\rho x)^\nu} = \frac{(1 - \rho)^{2(\alpha+1-\nu)}(1 + \rho)^{2(\beta+1-\nu)}}{2^{2(\alpha+\beta)+3-2\nu} \rho^{\alpha+\beta+2-\nu}} \times \sum_{n=0}^{\infty} \frac{(\alpha + \beta + 2n + 1)\Gamma(\alpha + \beta + n + 1)(\nu)_n}{\Gamma(\alpha + 1 + n)\Gamma(\beta + 1 + n)} Q_{n+\nu-1}^{(\alpha+1-\nu, \beta+1-\nu)} \left(\frac{1 + \rho^2}{2\rho} \right) P_n^{(\alpha, \beta)}(x).$$

Proof. Consider the generating function for Gegenbauer polynomials (see for instance, (18.12.4) in Olver *et al.* (2010))

$$\frac{1}{(1 + \rho^2 - 2\rho x)^\nu} = \sum_{n=0}^{\infty} \rho^n C_n^\nu(x) = \sum_{n=0}^{\infty} \rho^n \frac{(2\nu)_n}{(\nu + \frac{1}{2})_n} P_n^{(\nu-1/2, \nu-1/2)}(x), \quad (55)$$

where we have expressed the Gegenbauer polynomial as a symmetric Jacobi polynomial using (41). Now we utilize the connection relation for Jacobi polynomials (see for instance Ismail (2005) [14, p. 256]), namely

$$P_n^{(\gamma, \delta)}(x) = \sum_{k=0}^n c_{n,k}(\gamma, \delta; \alpha, \beta) P_k^{(\alpha, \beta)}(x),$$

where $\gamma, \delta > -1$ and

$$c_{n,k}(\gamma, \delta; \alpha, \beta) = \frac{(\gamma + k + 1)_{n-k}(n + \gamma + \delta + 1)_k \Gamma(\alpha + \beta + k + 1)}{(n - k)! \Gamma(\alpha + \beta + 2k + 1)} \times {}_3F_2 \left(\begin{matrix} -n + k, n + k + \gamma + \delta + 1, \alpha + k + 1 \\ \gamma + k + 1, \alpha + \beta + 2k + 2 \end{matrix} ; 1 \right).$$

The generalized hypergeometric function ${}_3F_2 : \mathbf{C}^3 \times (\mathbf{C} \setminus -\mathbf{N}_0)^2 \times \{z \in \mathbf{C} : |z| < 1\} \rightarrow \mathbf{C}$ can be defined as follows

$${}_3F_2 \left(\begin{matrix} a_1, a_2, a_3 \\ b_1, b_2 \end{matrix} ; z \right) := \sum_{n=0}^{\infty} \frac{(a_1)_n (a_2)_n (a_3)_n}{(b_1)_n (b_2)_n} \frac{z^n}{n!}. \quad (56)$$

Utilizing (56) in (55), reversing the order of the summations, and shifting the n index yields after some manipulations

$$\frac{1}{(1 + \rho^2 - 2\rho x)^\nu} = \frac{\sqrt{\pi}}{2^{2\nu-1} \Gamma(\nu)} \sum_{k=0}^{\infty} \frac{\rho^k \Gamma(2\nu + 2k) \Gamma(\alpha + \beta + k + 1)}{\Gamma(\nu + k + \frac{1}{2}) \Gamma(\alpha + \beta + 2k + 1)} P_k^{(\alpha, \beta)}(x) \times \sum_{n=0}^{\infty} \frac{\rho^n (2\nu + 2k)_n}{n!} {}_3F_2 \left(\begin{matrix} -n, n + 2\nu + 2k, \alpha + k + 1 \\ \nu + k + \frac{1}{2}, \alpha + \beta + 2k + 2 \end{matrix} ; 1 \right).$$

By substituting the definition of the ${}_3F_2$ generalized hypergeometric function (56) in the sum over n and reversing the orders of the resulting two summations, it follows that

$$\sum_{n=0}^{\infty} \frac{\rho^n (2\nu + 2k)_n}{n!} {}_3F_2 \left(\begin{matrix} -n, n + 2\nu + 2k, \alpha + k + 1 \\ \nu + k + \frac{1}{2}, \alpha + \beta + 2k + 2 \end{matrix} ; 1 \right)$$

$$= \frac{1}{(1-\rho)^{2\nu+2k}} {}_2F_1 \left(\begin{matrix} \nu+k, \alpha+k+1 \\ \alpha+\beta+2k+2 \end{matrix} ; \frac{-4\rho}{(1-\rho)^2} \right). \quad (57)$$

If we apply the right-hand side of (57) to the definition of the Jacobi function of the second kind $Q_\gamma^{(\alpha,\beta)} : \mathbf{C} \setminus (-\infty, 1] \rightarrow \mathbf{C}$ (cf. [10, (10.8.18)]), namely

$$Q_\gamma^{(\alpha,\beta)}(z) := \frac{2^{\alpha+\beta+\gamma} \Gamma(\alpha+\gamma+1) \Gamma(\beta+\gamma+1)}{\Gamma(\alpha+\beta+2\gamma+2)(z-1)^{\alpha+\gamma+1}(z+1)^\beta} {}_2F_1 \left(\begin{matrix} \gamma+1, \alpha+\gamma+1 \\ \alpha+\beta+2\gamma+2 \end{matrix} ; \frac{2}{1-z} \right), \quad (58)$$

where $\alpha+\gamma, \beta+\gamma \notin -\mathbf{N}$, noting Theorem 12.7.3 (Expansion of an analytic function in terms of orthogonal polynomials) in Szegő (1959) we obtain the desired result. ■

Lemma 5.2. *Let $n \in \mathbf{N}_0$, $\mu \in \mathbf{C} \setminus \{-\frac{1}{2}, -\frac{3}{2}, -\frac{5}{2}, \dots\}$, $\nu \in \mathbf{C} \setminus -\mathbf{N}_0$, $z \in \mathbf{C} \setminus (-\infty, 1]$. Then*

$$Q_{n+\nu-1}^{(\mu-\nu+1/2, \mu-\nu+1/2)}(z) = \frac{2^{\mu-\nu+1/2} \Gamma(\mu+n+1/2) e^{i\pi(\mu-\nu+1/2)}}{\Gamma(\nu+n)(z^2-1)^{(\mu-\nu)/2+1/4}} Q_{n+\mu-1/2}^{\nu-\mu-1/2}(z).$$

Proof. Comparison of the definition of Jacobi function of the second kind (58), with the definition of the associated Legendre function of the second kind $Q_\nu^\mu : \mathbf{C} \setminus (-\infty, 1] \rightarrow \mathbf{C}$ ([19, entry 24, p. 161])

$$Q_\nu^\mu(z) := \frac{e^{i\pi\mu} 2^\nu \Gamma(\nu+1) \Gamma(\nu+\mu+1) (z+1)^{\mu/2}}{\Gamma(2\nu+2)(z-1)^{b/2+a+1}} {}_2F_1 \left(\begin{matrix} \nu+1, \nu+\mu+1 \\ 2+2\nu \end{matrix} ; \frac{2}{1-z} \right),$$

and elsewhere in z by analytic continuation, completes the proof. ■

Corollary 5.3. *Let $\nu \in \mathbf{C} \setminus -\mathbf{N}_0$, with $\mu \in (-1/2, \infty) \setminus \{0\}$, and $z \in \mathbf{C} \setminus (-\infty, 1]$ on any ellipse with foci at ± 1 with x in the interior of that ellipse. Then*

$$\frac{1}{(z-x)^\nu} = \frac{2^{\mu+1/2} \Gamma(\mu) e^{i\pi(\mu-\nu+1/2)}}{\sqrt{\pi} \Gamma(\nu) (z^2-1)^{(\nu-\mu)/2-1/4}} \sum_{n=0}^{\infty} (n+\mu) Q_{n+\mu-1/2}^{\nu-\mu-1/2}(z) C_n^\mu(x). \quad (59)$$

If one substitutes $z = (1+\rho^2)/(2\rho)$ in (59) with $0 < |\rho| < 1$, then one obtains an alternate expression with $x \in [-1, 1]$, namely

$$\begin{aligned} \frac{1}{(1+\rho^2-2\rho x)^\nu} &= \frac{\Gamma(\mu) e^{i\pi(\mu-\nu+1/2)}}{\sqrt{\pi} \Gamma(\nu) \rho^{\mu+1/2} (1-\rho^2)^{\nu-\mu-1/2}} \\ &\quad \times \sum_{n=0}^{\infty} (n+\mu) Q_{n+\mu-1/2}^{\nu-\mu-1/2} \left(\frac{1+\rho^2}{2\rho} \right) C_n^\mu(x). \end{aligned}$$

Proof. Let $\alpha = \beta = \mu - 1/2$ in Theorem 5.1, use Lemma 5.2 and the definition of the Gegenbauer polynomial in terms of a symmetric Jacobi polynomial (41). ■

Corollary 5.4. *Let $d \in \{3, 4, 5, \dots\}$, $\nu \in \mathbf{C} \setminus \{0, 2, 4, \dots\}$, $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$ with $r = \|\mathbf{x}\|$, $r' = \|\mathbf{x}'\|$, and $\cos \gamma = (\mathbf{x}, \mathbf{x}')/(rr')$. Then*

$$\|\mathbf{x} - \mathbf{x}'\|^\nu = \frac{e^{i\pi(\nu+d-1)/2} \Gamma\left(\frac{d-2}{2}\right) (r_>^2 - r_<^2)^{(\nu+d-1)/2}}{2\sqrt{\pi} \Gamma\left(-\frac{\nu}{2}\right) (rr')^{(d-1)/2}} \times \sum_{n=0}^{\infty} (2n+d-2) Q_{n+(d-3)/2}^{(1-\nu-d)/2} \left(\frac{r^2 + r'^2}{2rr'}\right) C_n^{d/2-1}(\cos \gamma), \quad (60)$$

where $r_{\leq} = \min_{\max} \{r, r'\}$.

Proof. Map $\nu \mapsto -\nu/2$, and then directly substitute $z = (r^2 + r'^2)/(2rr')$, $x = \cos \gamma$ in (59), using $\|\mathbf{x} - \mathbf{x}'\| = \sqrt{2rr'}\sqrt{z-x}$, $\mathbf{x}, \mathbf{x}' \in \mathbf{R}^d$ for $d \geq 3$. ■

Corollary 5.5. *Let $\nu \in \mathbf{C} \setminus -\mathbf{N}_0$, $\mu \in (-1/2, \infty) \setminus \{0\}$, and $x, z \in \mathbf{C}$ such that $z \in \mathbf{C} \setminus (-\infty, 1]$ lie on any ellipse with foci at ± 1 with x in the interior of that ellipse. Then*

$$\frac{1}{(z-x)^\nu} = \frac{\sqrt{2} e^{-i\pi(\nu-1/2)}}{\sqrt{\pi} \Gamma(\nu) (z^2-1)^{\nu/2-1/4}} \sum_{n=0}^{\infty} \epsilon_n T_n(x) Q_{n-1/2}^{\nu-1/2}(z). \quad (61)$$

If one substitutes $z = (1 + \rho^2)/(2\rho)$ in (61) with $0 < |\rho| < 1$, then one obtains an alternate expression with $x \in [-1, 1]$, namely

$$\frac{1}{(1 + \rho^2 - 2\rho x)^\nu} = \frac{e^{i\pi(\nu-1/2)}}{\sqrt{\pi\rho} \Gamma(\nu) (1 - \rho^2)^{\nu-1/2}} \sum_{n=0}^{\infty} \epsilon_n T_n(x) Q_{n-1/2}^{\nu-1/2} \left(\frac{1 + \rho^2}{2\rho}\right).$$

Proof. Take the limit as $\mu \rightarrow 0$ on the right-hand side of (59) and use

$$\lim_{\mu \rightarrow 0} \frac{n + \mu}{\mu} C_n^\mu(x) = \epsilon_n T_n(x) \quad (62)$$

(see for instance (6.4.13) in Andrews, Askey & Roy (1999) [2]), where $T_n : \mathbf{C} \rightarrow \mathbf{C}$ is the Chebyshev polynomial of the first kind defined as (see section 5.7.2 in Magnus, Oberhettinger & Soni (1966) [19])

$$T_n(x) := {}_2F_1 \left(\begin{matrix} -n, n \\ \frac{1}{2} \end{matrix} ; \frac{1-x}{2} \right).$$

■

Note that (61) is given in Cohl & Dominici (2011) [5, (3.10)], so (54) and (59) represent generalizations of that formula.

6. Addition theorems in Vilenkin's polyspherical coordinates

As mentioned in the introduction, by comparing azimuthal Fourier expansions for fundamental solutions of the polyharmonic equation on $\mathbf{R}^d$ for $d \geq 3$, with eigenfunction expansions for these fundamental solutions in rotationally-invariant coordinate systems, one can construct multi-summation and multi-integration addition theorems.

We have already constructed some of these addition theorems on $\mathbf{R}^3$ (see Cohl *et al.* (2000) [8]; Cohl *et al.* (2001) [7]). In Cohl *et al.* (2000) [8] addition theorems are derived in this fashion for fundamental solutions of the Laplacian on $\mathbf{R}^3$, namely in cylindrical, oblate spheroidal, prolate spheroidal, parabolic, bispherical and toroidal coordinate systems. In Cohl *et al.* (2001) [7] an addition theorem is derived from a fundamental solution of Laplace's equation in type **ba** (standard) spherical coordinates (we generalize that addition theorem below (74)). In circular cylindrical and rotational parabolic coordinates, the derived addition theorems are given in terms of definite integrals. In oblate spheroidal, prolate spheroidal, bispherical, toroidal, and spherical coordinates, the derived addition theorems are given in terms of sums. As is indicated by the above presentation, one may construct addition theorems in this manner in any rotationally-invariant coordinate system which yields solutions through separation of variables for the polyharmonic equation. In a similar setting, addition theorems may be generated for other inhomogeneous linear partial differential equations, such as for the Helmholtz, wave and heat equations in arbitrary dimensions. Furthermore an extension of this concept is possible when working with linear partial differential operators on Riemannian manifolds, such as for the Laplace-Beltrami operator (see Cohl & Kalnins (2012) [6, section 5.1]). Once a Fourier expansion for a fundamental solution is obtained for a partial differential operator on a Riemannian manifold, one must construct eigenfunction expansions for a fundamental solution corresponding to that operator and identify those nested multi-summation and multi-integration eigenfunction expansions which correspond to the Fourier coefficient for that operator.

Suppose one adopts for $\mathbf{R}^d$ a transformation using standard polyspherical coordinates (35). New addition theorems can then be generated by re-arranging the relevant sums over the hyperspherical harmonics, for instance

$$\sum_{l_1=0}^{l_{1\max}} \sum_K = \sum_{m=-l_{1\max}}^{l_{1\max}} \sum_{l_{d-2}=|m|}^{l_{1\max}} \sum_{l_{d-3}=l_{d-2}}^{l_{1\max}} \cdots \sum_{l_2=l_3}^{l_{1\max}} \sum_{l_1=l_2}^{l_{1\max}}. \quad (63)$$

In (63), one must sum over all allowed quantum numbers $\{l_1\} \cup K$. This is accomplished by taking the limit on both sides of (63) as $l_{1\max} \rightarrow \infty$. The forward sum on the left-hand side of (63) is now to be interpreted in a reverse fashion as in the right-hand side. Take for instance the limit $l_{1\max} \rightarrow \infty$ in the final sum, i.e., $\{l_1 \in \mathbf{N}_0 : l_1 \geq l_2\}$. Once l_1 has been summed over and eliminated from the sum, then one must sum over the remaining quantum numbers. In this limit, one then sums over l_2 in the set $\{l_2 \in \mathbf{N}_0 : l_2 \geq l_3\}$, eliminating l_2 from the sum. One continues this process until we have eliminated the contribution from all quantum numbers except m and finally we are left with a series

which only depends on m . Since we have already identified the azimuthal Fourier coefficients in section 5 (the terms corresponding to m with the other quantum numbers completely summed over), then by equating these two expressions, we can construct multi-index addition theorems in any polyspherical coordinate system. In the discussion below, we give explicit examples of this procedure.

6.1. Power-law addition theorem on $\mathbf{R}^d$ for $d \geq 3$ in standard polyspherical coordinates

In standard polyspherical coordinates (35) we have the following addition theorem.

Theorem 6.1. *Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{Z}$, $\theta_i \in [0, \pi]$, for $1 \leq i \leq d-2$, $r, r' \in [0, \infty)$ with $d \geq 3$. Then*

$$\begin{aligned}
Q_{m-1/2}^{-(\nu+1)/2}(\chi) &= \sqrt{2} e^{i\pi(d-2)/2} \pi^{(d-4)/2} \left(2rr' \prod_{i=1}^{d-2} \sin \theta_i \sin \theta'_i \right)^{-\nu/2} \\
&\times (\chi^2 - 1)^{-(\nu+1)/4} \frac{(r_{>}^2 - r_{<}^2)^{(\nu+d-1)/2}}{(rr')^{(d-1)/2}} \\
&\times \sum_{l_{d-2}=m}^{\infty} \frac{(2l_{d-2}+1)(l_{d-2}-m)!}{(l_{d-2}+m)!} P_{l_{d-2}}^m(\cos \theta_{d-2}) P_{l_{d-2}}^m(\cos \theta'_{d-2}) \\
&\times \sum_{l_{d-3}=l_{d-2}}^{\infty} \Theta_{d-3}^d(l_{d-3}, l_{d-2}; \theta_{d-3}) \Theta_{d-3}^d(l_{d-3}, l_{d-2}; \theta'_{d-3}) \\
&\vdots \\
&\times \sum_{l_2=l_3}^{\infty} \Theta_2^d(l_2, l_3; \theta_2) \Theta_2^d(l_2, l_3; \theta'_2) \\
&\times \sum_{l_1=l_2}^{\infty} \Theta_1^d(l_1, l_2; \theta_1) \Theta_1^d(l_1, l_2; \theta'_1) Q_{l_1+(d-3)/2}^{(1-\nu-d)/2} \left(\frac{r^2 + r'^2}{2rr'} \right), \tag{64}
\end{aligned}$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \sum_{i=1}^{d-2} \cos \theta_i \cos \theta'_i \prod_{j=1}^{i-1} \sin \theta_j \sin \theta'_j}{2rr' \prod_{i=1}^{d-2} \sin \theta_i \sin \theta'_i}. \tag{65}$$

Proof. If we adopt standard polyspherical coordinates (35) (see figure 4), then one can obtain an eigenfunction expansion for a power-law fundamental solution of the polyharmonic equation in standard polyspherical coordinates using the Gegenbauer expansion (60) with the addition theorem for hyperspherical harmonics (1), (50). The Fourier expansion for a power-law fundamental solution of the polyharmonic equation

is obtained from the algebraic expression for a power-law fundamental solution of the polyharmonic equation (cf. (7)) and (61), namely

$$\begin{aligned} \|\mathbf{x} - \mathbf{x}'\|^\nu &= \sqrt{\frac{\pi}{2}} \frac{e^{i\pi(\nu+1)/2}}{\Gamma(-\nu/2)} \left(2rr' \prod_{i=1}^{d-2} \sin \theta_i \sin \theta'_i \right)^{\nu/2} (\chi^2 - 1)^{(\nu+1)/4} \\ &\quad \times \sum_{m=-\infty}^{\infty} e^{im(\phi-\phi')} Q_{m-1/2}^{-(\nu+1)/2}(\chi), \end{aligned} \quad (66)$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \sum_{i=1}^{d-2} \cos \theta_i \cos \theta'_i \prod_{j=1}^{i-1} \sin \theta_j \sin \theta'_j}{2rr' \prod_{i=1}^{d-2} \sin \theta_i \sin \theta'_i}. \quad (67)$$

The Gegenbauer expansion yields

$$\begin{aligned} \|\mathbf{x} - \mathbf{x}'\|^\nu &= \frac{4e^{i\pi(\nu+d-1)/2} \pi^{(d-1)/2} (r_{>}^2 - r_{<}^2)^{(\nu+d-1)/2}}{\Gamma(-\frac{\nu}{2}) (rr')^{(d-1)/2}} \\ &\quad \times \sum_{l_1, K} Q_{l_1+(d-3)/2}^{(1-\nu-d)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) Y_{l_1}^K(\hat{\mathbf{x}}) \overline{Y_{l_1}^K(\hat{\mathbf{x}}')}. \end{aligned} \quad (68)$$

Now if we expand the product of polyspherical harmonics in (68) with (50), (63), we obtain

$$\begin{aligned} \|\mathbf{x} - \mathbf{x}'\|^\nu &= \sum_{m=0}^{\infty} \cos(m(\phi - \phi')) \frac{\pi^{(d-3)/2} e^{i\pi(\nu+d-1)/2} \epsilon_m (r_{>}^2 - r_{<}^2)^{(\nu+d-1)/2}}{\Gamma(-\frac{\nu}{2}) (rr')^{(d-1)/2}} \\ &\quad \times \sum_{l_{d-2}=m}^{\infty} \frac{(2l_{d-2} + 1)(l_{d-2} - m)!}{(l_{d-2} + m)!} P_{l_{d-2}}^m(\cos \theta_{d-2}) P_{l_{d-2}}^m(\cos \theta'_{d-2}) \\ &\quad \times \sum_{l_{d-3}=l_{d-2}}^{\infty} \Theta_{d-3}^d(l_{d-3}, l_{d-2}; \theta_{d-3}) \Theta_{d-3}^d(l_{d-3}, l_{d-2}; \theta'_{d-3}) \\ &\quad \vdots \\ &\quad \times \sum_{l_2=l_3}^{\infty} \Theta_2^d(l_2, l_3; \theta_2) \Theta_2^d(l_2, l_3; \theta'_2) \\ &\quad \times \sum_{l_1=l_2}^{\infty} \Theta_1^d(l_1, l_2; \theta_1) \Theta_1^d(l_1, l_2; \theta'_1) Q_{l_1+(d-3)/2}^{(1-\nu-d)/2} \left(\frac{r^2 + r'^2}{2rr'} \right), \end{aligned} \quad (69)$$

where $\Theta_j^d(l_j, l_{j+1}; \theta)$, for $1 \leq j \leq d-2$ is defined in (52). By comparing the Fourier coefficients of (69) with (66), we complete the proof of this theorem. $\blacksquare$

This is just one example of a derived multi-summation addition theorem for arbitrary dimensions. There are an unlimited number of such straightforward examples to generate. In the next section we derive another example which is valid on $\mathbf{R}^d$ where d is given by a power of two, i.e., generalized Hopf coordinates (36).

6.2. Power-law addition theorem on $\mathbf{R}^{2^q}$ for $q \geq 2$ in generalized Hopf coordinates

Theorem 6.2. Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m_1 \in \mathbf{Z}$, $r, r' \in [0, \infty)$, $\vartheta_i \in [0, \frac{\pi}{2}]$ with $1 \leq i \leq 2^{q-1} - 1$, $\phi_i \in [0, 2\pi)$ such that $1 \leq i \leq 2^{q-1}$. Then

$$\begin{aligned}
Q_{m_1-1/2}^{-(\nu+1)/2}(\chi) &= \frac{-\left(\prod_{j=1}^{q-1} \cos \vartheta_{2^{j-1}} \cos \vartheta'_{2^{j-1}}\right)^{-\nu/2}}{2^{(\nu+1)/2}(\chi^2 - 1)^{(\nu+1)/4}} \left(\frac{r_{>}^2 - r_{<}^2}{rr'}\right)^{(\nu+2^q-1)/2} \\
&\times \sum_{m_2=0}^{\infty} \epsilon_{m_2} \cos(m_2(\phi_2 - \phi'_2)) \cdots \sum_{m_{2^{q-1}}=0}^{\infty} \epsilon_{m_{2^{q-1}}} \cos(m_{2^{q-1}}(\phi_{2^{q-1}} - \phi'_{2^{q-1}})) \\
&\times \sum_{n_{2^{q-1}-1}=0}^{\infty} \Upsilon_{2^{q-1}-1}^q \left(\begin{matrix} n_{2^{q-1}-1} \\ |m_{2^{q-1}-1}|, |m_{2^{q-1}}| \end{matrix} ; \vartheta_{2^{q-1}-1} \right) \Upsilon_{2^{q-1}-1}^q \left(\begin{matrix} n_{2^{q-1}-1} \\ |m_{2^{q-1}-1}|, |m_{2^{q-1}}| \end{matrix} ; \vartheta'_{2^{q-1}-1} \right) \\
&\vdots \\
&\times \sum_{n_{2^{q-2}}=0}^{\infty} \Upsilon_{2^{q-2}}^q \left(\begin{matrix} n_{2^{q-2}} \\ |m_1|, |m_2| \end{matrix} ; \vartheta_{2^{q-2}} \right) \Upsilon_{2^{q-2}}^q \left(\begin{matrix} n_{2^{q-2}} \\ |m_1|, |m_2| \end{matrix} ; \vartheta'_{2^{q-2}} \right) \\
&\times \sum_{n_{2^{q-2}-1}=0}^{\infty} \Upsilon_{2^{q-2}-1}^q \left(\begin{matrix} n_{2^{q-2}-1} \\ l_{2^{q-1}-2}, l_{2^{q-1}-1} \end{matrix} ; \vartheta_{2^{q-2}-1} \right) \Upsilon_{2^{q-2}-1}^q \left(\begin{matrix} n_{2^{q-2}-1} \\ l_{2^{q-1}-2}, l_{2^{q-1}-1} \end{matrix} ; \vartheta'_{2^{q-2}-1} \right) \\
&\vdots \\
&\times \sum_{n_1=0}^{\infty} \Upsilon_1^q \left(\begin{matrix} n_1 \\ l_2, l_3 \end{matrix} ; \vartheta_1 \right) \Upsilon_1^q \left(\begin{matrix} n_1 \\ l_2, l_3 \end{matrix} ; \vartheta'_1 \right) Q_{2n_1+l_2+l_3+(2^q-3)/2}^{(1-\nu-2^q)/2} \left(\frac{r^2 + r'^2}{2rr'} \right), \tag{70}
\end{aligned}$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \cos \gamma + 2rr' \cos(\phi_1 - \phi'_1) \prod_{j=1}^{q-1} \cos \vartheta_{2^{j-1}} \cos \vartheta'_{2^{j-1}}}{2rr' \prod_{j=1}^{q-1} \cos \vartheta_{2^{j-1}} \cos \vartheta'_{2^{j-1}}}. \tag{71}$$

Proof. If we adopt generalized Hopf coordinates (36) (see figure 5), then we can use the corresponding harmonics (53) in combination with the addition theorem for hyperspherical harmonics (1). We compare the Gegenbauer expansion for powers of the

distance (60) with the Fourier expansion

$$\|\mathbf{x} - \mathbf{x}'\|^\nu = \frac{\sqrt{2}ie^{i\pi\nu/2} \left(2rr' \prod_{j=1}^{q-1} \cos \vartheta_{2^j-1} \cos \vartheta'_{2^j-1} \right)^{\nu/2}}{\sqrt{\pi} \Gamma\left(\frac{-\nu}{2}\right) (\chi^2 - 1)^{-(\nu+1)/4}} \times \sum_{m_1=-\infty}^{\infty} e^{im_1(\phi_1 - \phi'_1)} Q_{m_1-1/2}^{-(\nu+1)/2}(\chi), \quad (72)$$

where $\chi > 1$ is given by (71). Notice that χ is independent of $\phi_1 - \phi'_1$. By using the Gegenbauer expansion (60) and inserting the appropriate Gegenbauer polynomial using the addition theorem for hyperspherical harmonics (1), we obtain

$$\|\mathbf{x} - \mathbf{x}'\|^\nu = \frac{-2i\pi^{2^{q-1}} e^{i\pi\nu/2} (r_{>}^2 - r_{<}^2)^{(\nu+2^q-1)/2}}{\sqrt{\pi} \Gamma\left(\frac{-\nu}{2}\right) (rr')^{(2^q-1)/2}} \times \sum_{l_1, K} Q_{l_1+(2^q-3)/2}^{(1-\nu-2^q)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) Y_{l_1}^K(\widehat{\mathbf{x}}) \overline{Y_{l_1}^K(\widehat{\mathbf{x}}')}. \quad (73)$$

By expanding the product of polyspherical harmonics in (73) with (53) expressed in terms of the surrogate quantum number n and reversing the order of the sums, we obtain a multi-summation expression for the power of the Euclidean distance between two points in generalized Hopf coordinates. Through comparison of the resulting equation with the m_1 Fourier coefficients of (72), we derive (70), a multi-summation addition theorem for the associated Legendre function of the second kind with argument χ (cf. (71)). $\blacksquare$

6.3. Power-law addition theorems on $\mathbf{R}^3$

In $d = 3$ there are two ways to construct polyspherical coordinates, with trees of type **b'a** and **ba**.

6.3.1. Type **ba** coordinates

Corollary 6.3. *Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{N}_0$, $\theta, \theta' \in [0, \pi]$, $r, r' \in [0, \infty)$. Then*

$$Q_{m-1/2}^{-(\nu+1)/2}(\chi) = i\sqrt{\pi} 2^{-(\nu+3)/2} (\sin \theta \sin \theta')^{-\nu/2} (\chi^2 - 1)^{-(\nu+1)/4} \left(\frac{r_{>}^2 - r_{<}^2}{rr'} \right)^{(\nu+2)/2} \times \sum_{l=m}^{\infty} (2l+1) \frac{(l-m)!}{(l+m)!} Q_l^{-(\nu+2)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) P_l^m(\cos \theta) P_l^m(\cos \theta'), \quad (74)$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \cos \theta \cos \theta'}{2rr' \sin \theta \sin \theta'}. \quad (75)$$

Proof. Taking $d = 3$ in (64) completes the proof. $\blacksquare$

Equation (74) is a generalization of one of the main results of Cohl *et al.* (2001) [7]. This can be observed if you substitute $\nu = -1$ in (74) (this corresponds to a fundamental solution of Laplace's equation on $\mathbf{R}^3$), then the associated Legendre function of the second kind on the right-hand side reduces to an elementary function through [1, (8.6.11)], producing

$$Q_{m-1/2}(\chi) = \pi \sqrt{\sin \theta \sin \theta'} \sum_{l=|m|}^{\infty} \frac{(l-m)!}{(l+m)!} \left(\frac{r_{\leq}}{r_{>}} \right)^{l+1/2} P_l^m(\cos \theta) P_l^m(\cos \theta'). \quad (76)$$

6.3.2. Type $\mathbf{b}'\mathbf{a}$ coordinates

Corollary 6.4. Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{Z}$, $\theta, \theta' \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, $r, r' \in [0, \infty)$. Then

$$Q_{m-1/2}^{-(\nu+1)/2}(\chi) = i\sqrt{\pi} 2^{-(\nu+3)/2} (\cos \theta \cos \theta')^{-\nu/2} (\chi^2 - 1)^{-(\nu+1)/4} \left(\frac{r_{>}^2 - r_{\leq}^2}{rr'} \right)^{(\nu+2)/2} \\ \times \sum_{l=|m|}^{\infty} (2l+1) \frac{(l-m)!}{(l+m)!} Q_l^{-(\nu+2)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) P_l^m(\sin \theta) P_l^m(\sin \theta'), \quad (77)$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \sin \theta \sin \theta'}{2rr' \cos \theta \cos \theta'}.$$

Proof. Using the method of trees, the map from type $\mathbf{ba}$ to type $\mathbf{b}'\mathbf{a}$ coordinates is given by $\theta, \theta' \mapsto \theta - \frac{\pi}{2}$. Under these transformations $\cos \theta, \cos \theta' \mapsto \sin \theta, \sin \theta'$ and $\sin \theta, \sin \theta' \mapsto -\cos \theta, -\cos \theta'$, as applied to (74). This completes the proof. $\blacksquare$

6.4. Power-law addition theorems on $\mathbf{R}^4$

In $d = 4$ there are five ways to construct polyspherical coordinates, with trees of type $\mathbf{b}^2\mathbf{a}$ (see figure 3a), $\mathbf{bb}'\mathbf{a}$ (see figure 3b), $\mathbf{b}'\mathbf{ba}$ (see figure 3c), $\mathbf{b}'^2\mathbf{a}$ (see figure 3d), $\mathbf{ca}^2$ (see figure 3e).

6.4.1. Type $\mathbf{b}^2\mathbf{a}$ coordinates

Corollary 6.5. Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{Z}$, $r, r' \in [0, \infty)$, $\theta_1, \theta'_1, \theta_2, \theta'_2 \in [0, \pi]$. Then

$$Q_{m-1/2}^{-(\nu+1)/2}(\chi) = \frac{-1}{2^{(\nu+1)/2}} \left(\frac{r_{>}^2 - r_{\leq}^2}{rr'} \right)^{(\nu+3)/2} (\chi^2 - 1)^{-(\nu+1)/4} (\sin \theta_1 \sin \theta'_1 \sin \theta_2 \sin \theta'_2)^{-\nu/2} \\ \times \sum_{l_2=|m|}^{\infty} \frac{2^{2l_2} (2l_2+1) (l_2!)^2 (l_2-m)!}{(l_2+m)!} (\sin \theta_1 \sin \theta'_1)^{l_2} P_{l_2}^m(\cos \theta_2) P_{l_2}^m(\cos \theta'_2)$$

$$\times \sum_{l_1=l_2}^{\infty} \frac{(l_1+1)(l_1-l_2)!}{(l_1+l_2+1)!} Q_{l_1+1/2}^{-(\nu+3)/2} \left(\frac{r^2+r'^2}{2rr'} \right) C_{l_1-l_2}^{l_2+1}(\cos \theta_1) C_{l_1-l_2}^{l_2+1}(\cos \theta'_1), \quad (78)$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \cos \theta_1 \cos \theta'_1 - 2rr' \sin \theta_1 \sin \theta'_1 \cos \theta_2 \cos \theta'_2}{2rr' \sin \theta_1 \sin \theta'_1 \sin \theta_2 \sin \theta'_2}. \quad (79)$$

Proof. Taking $d = 4$ in (64) completes the proof. ■

If you substitute $\nu = -2$ (a fundamental solution for the Laplacian on $\mathbf{R}^4$) in (78) then the Legendre functions of the second kind reduce to elementary functions through [1, (8.6.10-11)], and one obtains the following

$$\begin{aligned} \frac{(\chi^2 - 1)^{-1/2}}{(\chi + \sqrt{\chi^2 - 1})^m} &= 2 \sin \theta_1 \sin \theta'_1 \sin \theta_2 \sin \theta'_2 \\ &\times \sum_{l_2=|m|}^{\infty} \frac{2^{2l_2}(2l_2+1)(l_2!)^2(l_2-m)!}{(l_2+m)!} (\sin \theta_1 \sin \theta'_1)^{l_2} P_{l_2}^m(\cos \theta_2) P_{l_2}^m(\cos \theta'_2) \\ &\times \sum_{l_1=l_2}^{\infty} \frac{(l_1-l_2)!}{(l_1+l_2+1)!} \left(\frac{r_{\leq}}{r_{\geq}} \right)^{l_1+1} C_{l_1-l_2}^{l_2+1}(\cos \theta_1) C_{l_1-l_2}^{l_2+1}(\cos \theta'_1). \end{aligned}$$

6.4.2. Type $\mathbf{bb}'\mathbf{a}$ coordinates In type $\mathbf{bb}'\mathbf{a}$ coordinates, this is the addition theorem derived using the methods described above (see figure 3b)

Corollary 6.6. *Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{Z}$, $r, r' \in [0, \infty)$, $\theta_1, \theta'_1 \in [0, \pi]$, $\theta_2, \theta'_2 \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Then*

$$\begin{aligned} Q_{m-1/2}^{-(\nu+1)/2}(\chi) &= \frac{-1}{2^{(\nu+1)/2}} \left(\frac{r_{\geq}^2 - r_{\leq}^2}{rr'} \right)^{(\nu+3)/2} (\chi^2 - 1)^{-(\nu+1)/4} (\sin \theta_1 \sin \theta'_1 \cos \theta_2 \cos \theta'_2)^{-\nu/2} \\ &\times \sum_{l_2=|m|}^{\infty} \frac{2^{2l_2}(2l_2+1)(l_2!)^2(l_2-m)!}{(l_2+m)!} (\sin \theta_1 \sin \theta'_1)^{l_2} P_{l_2}^m(\sin \theta_2) P_{l_2}^m(\sin \theta'_2) \\ &\times \sum_{l_1=l_2}^{\infty} \frac{(l_1+1)(l_1-l_2)!}{(l_1+l_2+1)!} Q_{l_1+1/2}^{-(\nu+3)/2} \left(\frac{r^2+r'^2}{2rr'} \right) C_{l_1-l_2}^{l_2+1}(\cos \theta_1) C_{l_1-l_2}^{l_2+1}(\cos \theta'_1), \quad (80) \end{aligned}$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \cos \theta_1 \cos \theta'_1 - 2rr' \sin \theta_1 \sin \theta'_1 \sin \theta_2 \sin \theta'_2}{2rr' \sin \theta_1 \sin \theta'_1 \cos \theta_2 \cos \theta'_2}. \quad (81)$$

Proof. Using the method of trees, the map from type $\mathbf{b}^2\mathbf{a}$ to type $\mathbf{bb}'\mathbf{a}$ coordinates leaves θ_1 invariant and $\theta_2, \theta'_2 \mapsto \theta_2 - \frac{\pi}{2}, \theta'_2 - \frac{\pi}{2}$. Under these transformations $\cos \theta_2, \cos \theta'_2 \mapsto \sin \theta_2, \sin \theta'_2$ and $\sin \theta_2, \sin \theta'_2 \mapsto -\cos \theta_2, -\cos \theta'_2$, as applied to (80). This completes the proof. ■

6.4.3. Type $\mathbf{b}'\mathbf{ba}$ coordinates In type $\mathbf{b}'\mathbf{ba}$ coordinates, this is the addition theorem derived using the methods described above (see figure 3c)

Corollary 6.7. *Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{Z}$, $r, r' \in [0, \infty)$, $\theta_1, \theta'_1 \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, $\theta_2, \theta'_2 \in [0, \pi]$. Then*

$$\begin{aligned} Q_{m-1/2}^{-(\nu+1)/2}(\chi) &= \frac{-1}{2^{(\nu+1)/2}} \left(\frac{r_{>}^2 - r_{<}^2}{rr'} \right)^{(\nu+3)/2} (\chi^2 - 1)^{-(\nu+1)/4} (\cos \theta_1 \cos \theta'_1 \sin \theta_2 \sin \theta'_2)^{-\nu/2} \\ &\times \sum_{l_2=|m|}^{\infty} \frac{2^{2l_2} (2l_2+1) (l_2!)^2 (l_2-m)!}{(l_2+m)!} (\cos \theta_1 \cos \theta'_1)^{l_2} P_{l_2}^m(\cos \theta_2) P_{l_2}^m(\cos \theta'_2) \\ &\times \sum_{l_1=l_2}^{\infty} \frac{(l_1+1)(l_1-l_2)!}{(l_1+l_2+1)!} Q_{l_1+1/2}^{-(\nu+3)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) C_{l_1-l_2}^{l_2+1}(\sin \theta_1) C_{l_1-l_2}^{l_2+1}(\sin \theta'_1), \end{aligned} \quad (82)$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \sin \theta_1 \sin \theta'_1 - 2rr' \cos \theta_1 \cos \theta'_1 \cos \theta_2 \cos \theta'_2}{2rr' \cos \theta_1 \cos \theta'_1 \sin \theta_2 \sin \theta'_2}. \quad (83)$$

Proof. Using the method of trees, the map from type $\mathbf{b}^2\mathbf{a}$ to type $\mathbf{b}'\mathbf{ba}$ coordinates leaves θ_2 invariant and $\theta_1, \theta'_1 \mapsto \theta_1 - \frac{\pi}{2}, \theta'_1 - \frac{\pi}{2}$. Under these transformations $\cos \theta_1, \cos \theta'_1 \mapsto \sin \theta_1, \sin \theta'_1$ and $\sin \theta_1, \sin \theta'_1 \mapsto -\cos \theta_1, -\cos \theta'_1$, as applied to (80). This completes the proof. $\blacksquare$

6.4.4. Type $\mathbf{b}'^2\mathbf{a}$ coordinates In type $\mathbf{b}'^2\mathbf{a}$ coordinates, this is the addition theorem derived using the methods described above (see figure 3d)

Corollary 6.8. *Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m \in \mathbf{Z}$, $r, r' \in [0, \infty)$, $\theta_1, \theta'_1, \theta_2, \theta'_2 \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. Then*

$$\begin{aligned} Q_{m-1/2}^{-(\nu+1)/2}(\chi) &= \frac{-1}{2^{(\nu+1)/2}} \left(\frac{r_{>}^2 - r_{<}^2}{rr'} \right)^{(\nu+3)/2} (\chi^2 - 1)^{-(\nu+1)/4} (\cos \theta_1 \cos \theta'_1 \cos \theta_2 \cos \theta'_2)^{-\nu/2} \\ &\times \sum_{l_2=|m|}^{\infty} \frac{2^{2l_2} (2l_2+1) (l_2!)^2 (l_2-m)!}{(l_2+m)!} (\cos \theta_1 \cos \theta'_1)^{l_2} P_{l_2}^m(\sin \theta_2) P_{l_2}^m(\sin \theta'_2) \\ &\times \sum_{l_1=l_2}^{\infty} \frac{(l_1+1)(l_1-l_2)!}{(l_1+l_2+1)!} Q_{l_1+1/2}^{-(\nu+3)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) C_{l_1-l_2}^{l_2+1}(\sin \theta_1) C_{l_1-l_2}^{l_2+1}(\sin \theta'_1), \end{aligned} \quad (84)$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \sin \theta_1 \sin \theta'_1 - 2rr' \cos \theta_1 \cos \theta'_1 \sin \theta_2 \sin \theta'_2}{2rr' \cos \theta_1 \cos \theta'_1 \cos \theta_2 \cos \theta'_2}. \quad (85)$$

Proof. Using the method of trees, the map from type $\mathbf{b}^2\mathbf{a}$ to type $\mathbf{b}'^2\mathbf{a}$ coordinates $\theta_1, \theta'_1 \mapsto \theta_1 - \frac{\pi}{2}, \theta'_1 - \frac{\pi}{2}$ and $\theta_2, \theta'_2 \mapsto \theta_2 - \frac{\pi}{2}, \theta'_2 - \frac{\pi}{2}$. Under these transformations $\cos \theta_1, \cos \theta'_1, \cos \theta_2, \cos \theta'_2 \mapsto \sin \theta_1, \sin \theta'_1, \sin \theta_2, \sin \theta'_2$ and $\sin \theta_1, \sin \theta'_1 \mapsto -\cos \theta_1, -\cos \theta'_1, -\cos \theta_2, -\cos \theta'_2$, as applied to (80). This completes the proof. $\blacksquare$

6.4.5. *Type $\mathbf{ca}^2$ coordinates* In type $\mathbf{ca}^2$ coordinates, this is the addition theorem derived using the methods described above

Corollary 6.9. *Let $\nu \in \mathbf{C} \setminus \{2m, 2m+2, 2m+4, \dots\}$, $m_1 \in \mathbf{Z}$, $r, r' \in [0, \infty)$, $\vartheta, \vartheta' \in [0, \frac{\pi}{2}]$, $\phi_2, \phi'_2 \in [0, 2\pi)$. Then*

$$\begin{aligned}
Q_{m_1-1/2}^{-(\nu+1)/2}(\chi) &= \frac{-2^{-(\nu+1)/2}}{(\chi^2 - 1)^{(\nu+1)/4}} \left(\frac{r_{>}^2 - r_{<}^2}{rr'} \right)^{(\nu+3)/2} (\cos \vartheta \cos \vartheta')^{|m_1|-\nu/2} \\
&\times \sum_{m_2=0}^{\infty} \epsilon_{m_2} \cos(m_2(\phi_2 - \phi'_2)) (\sin \vartheta \sin \vartheta')^{m_2} \\
&\times \sum_{n=0}^{\infty} \frac{(2n + |m_1| + m_2 + 1)(|m_1| + m_2 + n)!n!}{(|m_1| + n)!(m_2 + n)!} \\
&\times Q_{2n+|m_1|+m_2+1/2}^{-(\nu+3)/2} \left(\frac{r^2 + r'^2}{2rr'} \right) P_n^{(m_2, |m_1|)}(\cos 2\vartheta) P_n^{(m_2, |m_1|)}(\cos 2\vartheta'), \quad (86)
\end{aligned}$$

where

$$\chi = \frac{r^2 + r'^2 - 2rr' \sin \vartheta \sin \vartheta' \cos(\phi_2 - \phi'_2)}{2rr' \cos \vartheta \cos \vartheta'}. \quad (87)$$

Proof. Taking $q = 2$ in (70) completes the proof. ■

If you substitute $\nu = -2$ in (86), then the Legendre functions of the second kind reduce to elementary functions through [1, (8.6.10-11)], and one obtains the following

$$\begin{aligned}
\frac{(\chi^2 - 1)^{-1/2}}{(\chi + \sqrt{\chi^2 - 1})^{m_1}} &= 2 (\cos \vartheta \cos \vartheta')^{|m_1|+1} \sum_{m_2=0}^{\infty} \epsilon_{m_2} \cos [m_2(\phi_2 - \phi'_2)] (\sin \vartheta \sin \vartheta')^{|m_2|} \\
&\times \sum_{n=0}^{\infty} \frac{(|m_1| + |m_2| + n)!n!}{(|m_1| + n)!(|m_2| + n)!} \left(\frac{r_{<}}{r_{>}} \right)^{|m_1|+|m_2|+2n+1} \\
&\times P_n^{(|m_2|, |m_1|)}(\cos 2\vartheta) P_n^{(|m_2|, |m_1|)}(\cos 2\vartheta'). \quad (88)
\end{aligned}$$

Note that in the addition theorems (86) and (88), that if you make the map $\vartheta \mapsto \vartheta - \frac{\pi}{2}$, then this transformation preserves the addition theorems such that $m_1 \leftrightarrow m_2$. This transformation is equivalent to swapping the position of ϕ_1 and ϕ_2 for the tree in figure 3e.

Acknowledgements

I would like to thank Tom ter Elst and Heather Macbeth for valuable discussions. Part of this work was conducted while H. S. Cohl was a National Research Council Research Postdoctoral Associate in the Information Technology Laboratory at the National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.

References

- [1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. U.S. Government Printing Office, Washington, D.C., 1972.
- [2] G. E. Andrews, R. Askey, and R. Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.
- [3] J. B. Boyling. Green’s functions for polynomials in the Laplacian. *Zeitschrift für Angewandte Mathematik und Physik*, 47(3):485–492, 1996.
- [4] H. S. Cohl. *Fourier and Gegenbauer expansions for fundamental solutions of the Laplacian and powers in $\mathbf{R}^d$ and $\mathbf{H}^d$* . PhD thesis, The University of Auckland, 2010. xiv+190 pages.
- [5] H. S. Cohl and D. E. Dominici. Generalized Heine’s identity for complex Fourier series of binomials. *Proceedings of the Royal Society A*, 467:333–345, 2011.
- [6] H. S. Cohl and E. G. Kalnins. Fourier and Gegenbauer expansions for a fundamental solution of the Laplacian in the hyperboloid model of hyperbolic geometry. *Journal of Physics A: Mathematical and Theoretical*, 45(14):145206, 2012.
- [7] H. S. Cohl, A. R. P. Rau, J. E. Tohline, D. A. Browne, J. E. Cazes, and E. I. Barnes. Useful alternative to the multipole expansion of $1/r$ potentials. *Physical Review A: Atomic and Molecular Physics and Dynamics*, 64(5):052509, Oct 2001.
- [8] H. S. Cohl, J. E. Tohline, A. R. P. Rau, and H. M. Srivastava. Developments in determining the gravitational potential using toroidal functions. *Astronomische Nachrichten*, 321(5/6):363–372, 2000.
- [9] T. H. Cormen, C. E. Leiserson, R. L. Rivest, and C. Stein. *Introduction to algorithms*. MIT Press, Cambridge, MA, second edition, 2001.
- [10] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Higher transcendental functions. Vol. II*. Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981.
- [11] U. Fano and A. R. P. Rau. *Symmetries in quantum physics*. Academic Press Inc., San Diego, CA, 1996.
- [12] I. M. Gel’fand and G. E. Shilov. *Generalized functions. Vol. 1*. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1964 [1977]. Properties and operations, Translated from the Russian by Eugene Saletan.
- [13] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Elsevier/Academic Press, Amsterdam, seventh edition, 2007. Translated from the Russian, Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger, With one CD-ROM (Windows, Macintosh and UNIX).
- [14] M. E. H. Ismail. *Classical and quantum orthogonal polynomials in one variable*, volume 98 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2005. With two chapters by Walter Van Assche, With a foreword by Richard A. Askey.
- [15] A. A. Izmet’shev, G. S. Pogosyan, A. N. Sissakian, and P. Winternitz. Contractions of Lie algebras and separation of variables. The n -dimensional sphere. *Journal of Mathematical Physics*, 40(3):1549–1573, 1999.
- [16] A. A. Izmet’shev, G. S. Pogosyan, A. N. Sissakian, and P. Winternitz. Contractions of Lie algebras and the separation of variables: interbase expansions. *Journal of Physics A: Mathematical and General*, 34(3):521–554, 2001.
- [17] E. G. Kalnins. *Separation of variables for Riemannian spaces of constant curvature*, volume 28 of *Pitman Monographs and Surveys in Pure and Applied Mathematics*. Longman Scientific & Technical, Harlow, 1986.
- [18] M. S. Kil’dyushov. Hyperspherical functions of the “tree” type in the n -body problem. *Soviet Journal of Nuclear Physics*, 15(1):113–118, 1972.
- [19] W. Magnus, F. Oberhettinger, and R. P. Soni. *Formulas and theorems for the special functions of mathematical physics*. Third enlarged edition. Die Grundlehren der mathematischen

- Wissenschaften, Band 52. Springer-Verlag New York, Inc., New York, 1966.
- [20] W. Miller, Jr. *Symmetry and separation of variables*. Addison-Wesley Publishing Co., Reading, Mass.-London-Amsterdam, 1977. With a foreword by Richard Askey, *Encyclopedia of Mathematics and its Applications*, Vol. 4.
 - [21] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST handbook of mathematical functions*. Cambridge University Press, Cambridge, 2010.
 - [22] L. Schwartz. *Théorie des distributions. Tome I*. Actualités Sci. Ind., no. 1091 = Publ. Inst. Math. Univ. Strasbourg 9. Hermann, Paris, 1950.
 - [23] N. J. A. Sloane and S. Plouffe. *The encyclopedia of integer sequences*. Academic Press Inc., San Diego, CA, 1995. With a separately available computer disk.
 - [24] R. P. Stanley. *Enumerative combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin.
 - [25] N. Ja. Vilenkin. *Special functions and the theory of group representations*. Translated from the Russian by V. N. Singh. Translations of Mathematical Monographs, Vol. 22. American Mathematical Society, Providence, R. I., 1968.
 - [26] N. Ja. Vilenkin, G. I. Kuznetsov, and Ja. A. Smorodinskii. Eigenfunctions of the Laplace operator providing representations of the $U(2)$, $SU(2)$, $SO(3)$, $U(3)$ and $SU(3)$ groups and the symbolic method. *Soviet Journal of Nuclear Physics*, 2:645–652, 1965.
 - [27] Z. Y. Wen and J. Avery. Some properties of hyperspherical harmonics. *Journal of Mathematical Physics*, 26(3):396–403, 1985.